



Dehydrated Culture Media  
Bases / Media Supplements

## Technical Information

### Rappaport Vassiliadis Medium, Modified

#### Product Code: DM 1880F

**Application:** - Rappaport Vassiliadis Medium, Modified is recommended for enrichment of *Salmonellae* in accordance with FDA BAM, 1998.

#### Composition\*\*

Ingredients	Gms / 1110 ml
Tryptone	5.000
Sodium chloride	8.000
Potassium dihydrogen phosphate	1.600
Magnesium chloride hexahydrate	18.73
Malachite green oxalate	0.040
Final pH ( at 25°C)	5.5±0.2

\*\*Formula adjusted, standardized to suit performance parameters

#### Principle & Interpretation

Rappaport Vassiliadis Medium, Modified is a modification of the formulation by Van Schothorst et al (1) and is recommended for the selective enrichment of *Salmonellae* from food specimens by FDA BAM, 1998(2). *Salmonella* generally survive at little high osmotic pressure, grow at slightly low pH and are resistant to malachite green compared to other bacteria. *Salmonellae* constitute the most taxonomically complex group of bacteria among the *Enterobacteriaceae* (3). Human *Salmonella* infections are most commonly caused by ingestion of food, water or milk contaminated by human or animal excreta. Contaminated eggs or foods containing eggs have also been a source of food borne salmonellosis.

Inoculate the samples into appropriate enriched media and incubate for 24 hrs for  $24 \pm 2$  h at 35°C. Transfer 0.1 ml mixture to 10 ml Rappaport Vassiliadis Medium, Modified (DM1880F) and another 1 ml into 10 ml Tetrathionate (TT) broth (DM1880F). Vortex and incubate at optimum temperature for  $24 \pm 2$ h depending upon the microbial load and type of the sample. These are further subcultured into XLD Agar or HE Agar, incubate the plates for  $24 \pm 2$ h at 35°C and observe for the appearance of typical salmonellae colonies. Blue-green to blue colonies will be appeared in XLD Agar and pink colonies with or without black centers on HE Agar.

The medium contains Tryptone which provides essential growth nutrients. Magnesium chloride raises the osmotic pressure in the medium. Malachite green is inhibitory to organisms other than *Salmonellae*. The low pH of the medium, combined with the presence of malachite green and magnesium chloride, helps to select for the highly resistant *Salmonella* species. Potassium phosphate buffers the medium to maintain the constant pH. Sodium chloride maintains the osmotic balance.

#### Methodology

Suspend 33.37 grams of dehydrated powder media in 1000 ml distilled water. Mix thoroughly & heat if necessary to dissolve the medium completely. Dispense as desired into tubes and sterilize by autoclaving at 10 lbs pressure (115°C) for 15 minutes.

#### Quality Control

##### Appearance

Light yellow to light blue homogeneous free flowing powder.



**Colour and Clarity**

Bluish green coloured, clear to slightly opalescent solution with slight precipitate

**Reaction**

Reaction of 3.34% w/v aqueous solution at 25°C. pH : 5.5±0.2

**pH Range**

5.30-5.70

**Cultural Response**

DM1880F: Cultural characteristics observed after an incubation at 42-43°C for 18-24 hours. After incubation, subculture on selective agar media like MacConkey Agar (DM1081) or XLD Agar (DM1031) and incubate at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth at 42±1°C	Recovery	Colour of Colony
<i>Escherichia coli</i> ATCC 25922	50-100	none-poor	<=10%	pink-red
<i>Salmonella Enteritidis</i> ATCC 13076	50-100	good-luxuriant	>=50%	colourless
<i>Salmonella Typhi</i> ATCC 6539	50-100	good-luxuriant	>=50%	colourless
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	good-luxuriant	>=50%	colourless

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and the prepared medium at 2 -8°C. Use before expiry date on the label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

## Further Reading

1. Van Schothorst, M., Renauld, A. and VanBeek, C 1987. Food Microbiol, 4.
2. FDA, U.S. 1998. Bacteriological Analytical Manual. 8 ed. Gaithersburg, MD: AOAC International.
3. Tindall, B. J., Crimont, P. A. D., Gorry, G. M. and Euzesny, B. P 2005. Int. J. Sys. Evol. Microbiol., 55.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performances parameters.