

Technical Information

Costein's LDS Test Medium

Product Code: DM 2621

Application: - Costein's LDS Test Medium is used for the identification of members of *Enterobacteriaceae* on the basis of lysine decarboxylase and hydrogen sulphide production.

Composition**

Ingredients	Gms / Litre
Meat peptone	4.500
Papaic digest of soyabean meal	2.000
Yeast extract	3.000
Sodium chloride	5.000
D-Glucose	1.000
L-Lysine monohydrochloride	10.000
Sodium thiosulphate	0.200
Ammonium iron (II) sulphate	0.200
Bromocresol purple	0.032
Agar	6.000
Final pH (at 25°C)	5.6±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Costein's LDS (Lysine Decarboxylase Sulfhydrase) Test Medium is formulated by Costein (1). This is the test culture medium for the simultaneous detection of lysine decarboxylase (LDC) and hydrogen sulphide production (2). Pietzsch used this medium on account of its high degree of reliability (3).

Meat peptone, papaic digest of soyabean meal and yeast extract in the medium supply nitrogen and other nutrients necessary to support bacterial growth. Sodium chloride helps to maintaining osmotic balance. Bromocresol purple is the pH indicator. Sodium thiosulphate serves as a reducing agent and helps to maintain a low oxygen tension in the medium. *Enterobacteriaceae* grows poorly at low pH; therefore, their growth is poor on this medium due to its low pH value of 5.6. LDC-positive species tend to neutralize the medium as a result of cadaverine production due to decarboxylation of lysine; the conditions for growth are thus improved and the pH indicator changes its colour from yellow to violet. Species which can also reduce thiosulphate to hydrogen sulphide, cause an additional blackening of the violet medium due to the precipitation of iron sulphide. LDC-negative species do not increase the pH value of the medium; the pH indicator does not undergo a colour change. Therefore growth of these microorganisms is poor and H₂S positive species are thus unable to produce any hydrogen sulphide e.g. *Citrobacter*, *Proteus vulgaris*, *Proteus mirabilis*, *Providentia*, *Enterobacter*, *Shigella* show yellow coloured growth. LDC-positive and H₂S positive organisms give black colour and may be surrounded by a violet zone e.g. *Salmonella*, *Edwardsiella* and others LDC-positive and H₂S negative organism are violet in colour e.g. *Escherichia*, *Klebsiella*, *Hafnia*, *Serratia* and some rare *Salmonella* and others.

Methodology

Suspend 32 grams of dehydrated media in 1000 ml distilled water. Mix thoroughly & dispense into test tubes to a depth of approximately 5 cm, if possible cover with a 5 mm layer of viscous paraffin. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Allow the tubes to solidify in a vertical position.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 0.6 % Agar gel

Colour and Clarity

Yellow coloured, clear to slightly opalescent gel forms in tubes as slants

Reaction

Reaction of 3.2% w/v aqueous solution at 25°C. pH : 5.6±0.2

pH Range

5.40-5.80

Cultural Response

DM2621: Cultural characteristics observed after an incubation at 35-37°C for 24-48 hours.

Organism	Inoculum (CFU)	Growth	Colour change of medium
Cultural Response			
<i>Escherichia coli</i> ATCC 25922	50-100	good	violet
<i>Serratia marcescens</i> ATCC 14756	50-100	good	violet
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	good	violet and black
<i>Salmonella Enteritidis</i> ATCC 13076	50-100	good	violet and black
<i>Salmonella Arizonae</i> ATCC 13314	50-100	good	violet and black
<i>Shigella flexneri</i> ATCC 12022	50-100	fair	yellow
<i>Citrobacter freundii</i> ATCC 8090	50-100	fair	yellow
<i>Proteus mirabilis</i> ATCC 29906	50-100	fair	yellow
<i>Proteus vulgaris</i> ATCC 13315	50-100	fair	yellow

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and the prepared medium at 2 - 8°C. Use before expiry date on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Costein I. D., 1968, Einzelnahrboden fur die biochemische Ausscheidung van Salmonella-und Arizona-Kulturen- Zbl. F. Bakt. I. Orig., 206:390-395.
2. Edwards P. R., Fife M. A., 1961, Lysine-Iron Agar in the detection of Arizona cultures, Appl. Microbiol., 9:478-480.
3. Pietzch O., 1975, Der Voges-Proskauer- Schnelltest und der Lysindecaboxylase-Sulphhydrase-Test, Zwei Schnellmethoden fur die Enterobacteriaceae Diagnostik., - Arch. lebensmittelhyg., 26: 23-24.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performances parameters.