

Technical Information

Fraser Broth Base

Product Code: DM 2327

Application: - Fraser Broth Base with added supplement is used as a primary as well as secondary enrichment medium, for the isolation and enumeration of *Listeria monocytogenes* from food and animal feeds.

Composition**

Ingredients	Gms / Litre
Peptic digest of animal tissue	5.000
Casein enzymic hydrolysate	5.000
Yeast extract	5.000
Meat extract	5.000
Sodium chloride	20.000
Disodium hydrogen phosphate.2H ₂ O	12.000
Potassium dihydrogen phosphate	1.350
Esculin	1.000
Lithium chloride	3.000
Final pH (at 25°C)	7.2±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Fraser Broth Base is based on the formulation of Fraser and Sperber (4) is used for the detection of *Listeria* species in food products. Fraser Broth Base is formulated so as to provide optimum conditions for the growth of *Listeria*. *L.monocytogenes* primarily causes meningitis, encephalitis or septicemia in humans (1,2). In pregnant women, *L. monocytogenes* often causes influenza like bacteremic illness that, if untreated, may lead to amnionitis and infection of the fetus, resulting in abortion, still birth or premature birth. Contaminated foods are the primary vehicles of transmission (3).

Peptic digest of animal tissue, casein enzymic hydrolysate, yeast extract, and meat extract make the media highly nutritive by supplying essential nutrients including carbonaceous and nitrogenous substances for the test organism. Phosphates maintain the buffers the medium. All *Listeria* species exhibit beta-glucosidase activity which is evident by the blackening of the media. *Listeria* species hydrolyze esculin (substituted glucoside) to glucose and esculetin. The latter combines with ferric ions of ferric ammonium citrate (MS 2141), resulting in the formation of 6-7 dihydroxycoumarin, a black brown complex. Ferric ammonium citrate also enhances the growth of *L.monocytogenes* (5). The high salt tolerance (of sodium chloride) of *Listeria* is used as means to inhibit the growth of Enterococci. Lithium chloride is also used to inhibit Enterococci, which also possess the ability to hydrolyze esculin. Growth of accompanying bacteria is largely prevented by the addition of Nalidixic acid and Acriflavin hydrochloride (MS 2125I).

The test sample under study is inoculated into the primary enrichment medium. After an incubation at 30°C for 18-24 hours, 0.1 ml is inoculated into Fraser Broth Base (DM 2327). After an incubation at 35-37°C for 24-48 hours, it is subcultured on *Listeria* Oxford Medium Base (DM 2145) or *Listeria* Identification Agar Base (PALCAM) DM 21064).

Methodology

Suspend 57.35 grams of dehydrated media in 1000 ml distilled water. Mix thoroughly & heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Aseptically add rehydrated contents of 1 vial of Fraser Selective Supplement (MS 2125I) and 2 vials of Fraser Supplement (MS 2141) to 1000 ml medium for primary enrichment or 1 vial of each to 500 ml medium for secondary enrichment. Shake well and dispense as desired.

Warning: Lithium chloride is harmful. Avoid bodily contact and inhalation of vapours. On contact with skin wash with plenty of water immediately.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity

Basal medium: Yellow coloured clear solution with slight precipitate. After addition: Fluorescent yellow coloured clear solution with slight precipitate forms in tubes.

Reaction

Reaction of 5.49% w/v aqueous solution at 25°C. pH : 7.2±0.2

pH Range

7.00-7.40

Cultural Response

DM 2327: Cultural characteristics observed on addition of MS 2125I and MS 2141 after an incubation at 35-37°C for 24-48 hours.

Organism	Inoculum (CFU)	Growth	Esculin Hydrolysis
Cultural Response			
<i>Escherichia coli</i> ATCC 25922	$\geq 10^3$	inhibited	-
<i>Enterococcus faecalis</i> ATCC 29212	$\geq 10^3$	inhibited	-
<i>Listeria monocytogenes</i> ATCC 19111	50-100	good-luxuriant	positive reaction, blackening of medium
<i>Listeria monocytogenes</i> ATCC 19112	50-100	good-luxuriant	positive reaction, blackening of medium
<i>Listeria monocytogenes</i> ATCC 19117	50-100	good-luxuriant	positive reaction, blackening of medium
<i>Listeria monocytogenes</i> ATCC 19118	50-100	good-luxuriant	positive reaction, blackening of medium
<i>Staphylococcus aureus</i> ATCC 25923	$\geq 10^3$	inhibited	-

Storage and Shelf Life

Dried Media: Store below 10-30°C in tightly closed container and prepared medium at 2-8°C. Use before expiry period on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Nieman R. E., and Lorber B., 1980, Rev. Infect. Dis. 2 : 207-227
2. Schuchat A. B., Swaminathan and C. V. Broome, Clin. Microbiol., Rev. 4 : 169-183
3. Murray P. R., Baron E. J., Jorgensen J. H., Pfaller M. A., Tenover F. C., Tenover F. C., (Eds.), 8th Ed., 2003, Manual of Clinical Microbiology, ASM, Washington, D.C.
4. Fraser and Sperber, 1988, J. Food Prot., 51:762-765.
5. Cowart R. E. and Foster B. G., 1985, J. Infect. Dis., 151:172.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.