

Dehydrated Culture Media Bases / Media Supplements

Technical Information

EC Blue Broth

Product Code: DM 2768

Application: - EC Blue Broth is used for detection and confirmation of *Escherichia coli* and total coliforms from water samples, using a combination of chromogenic and fluorogenic substrates.

Composition**					
Ingredients	Gms / Litre	Gms / Litre			
Peptone	5.000				
Sodium chloride	5.000				
Sodium pyruvate	1.000				
Potassium dihydrogen phosphate	1.000				
Dipotassium hydrogen phosphate	4.000				
Potassium nitrate	1.000				
Sodium lauryl sulphate	0.100				
IPTG	0.100				
X-Gal	0.100				
MUG	0.100				
Final pH (at 25°C)	7.1±0.2				
**Formula adjusted standardized to suit performance parameters					

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

EC Blue Broth is used for detection and confirmation of *Escherichia coli* and other coliforms. *Escherichia coli* can be distinguished from other coliforms by its unique ability to fluoresce in the presence of fluorogenic substrate (1, 2). The fluorogenic substrate, MUG is split by enzyme beta glucuronidase especially present in *Escherichia coli*. The reaction is indicated by the development of a blue fluorescence under UV light. The presence of total coliforms is indicated by blue- green colourations due to the cleavage of the chromogenic substrate (X-Gal). IPTG amplifies enzyme synthesis and increases the activity of beta -galactosidase.

Peptone supplies essential growth nutrients and is useful for the simultaneous detection of indole production. The phosphate salts provide buffering action for rapid growth of coliforms. Sodium chloride helps to maintain the osmotic balance. Sodium lauryl sulphate makes the medium selective by inhibiting accompanying microflora, especially the gram-positive organisms.

Methodology

Suspend 17.40 grams of dehydrated media in 1000 ml distilled water. Mix thoroughly & heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well before dispensing as desired.

Quality Control

Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity

Cream coloured clear solution having slight precipitate in tubes

Reaction

Reaction of 1.74 % w/v aqueous solution at 25°C. pH : 7.1±0.2

pH Range 6.90-7.30





Dehydrated Culture Media Bases / Media Supplements

Cultural Response

DM2768: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth	Colour change in medium	Fluorescence under UV light
Cultural Response Escherichia coli ATCC 25922	50-100	luxuriant	Bluish green	positive reaction
Klebsiella pneumoniae ATCC 13883	50-100	luxuriant	blue	negative reaction
Enterobacter aerogenes ATCC 13048	50-100	luxuriant	blue	negative reaction
Citrobacter freundii ATCC 8090	50-100	luxuriant	Bluish green	negative reaction
Salmonella Typhimurium ATCC 14028	50-100	luxuriant	colourless	negative reaction

Storage and Shelf Life

Dried Media: Store dehydrated and prepared medium at 2-8°C in tightly closed container. Use before expiry date on the label. **Prepared Media**: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Feng P.C.S. and Hartman P.A. ,1982, J.Appl. Environmental Microbiol. 43. 1320-1323. 2. Harsen W., and Yourassowsky, 1984, J.Clin. Microbiol.20.1177-1179.

Disclaimer:

• User must ensure suitability of the product(s) in their application prior to use.

• The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate

• Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.

• Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.

• Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performances parameters.

