

Technical Information

Kanamycin Esculin Azide MiVeg Broth Base

Product Code : VM1776A

Application:- Kanamycin Esculin Azide MiVeg Broth Base is used for isolation of *Enterococci* in foodstuffs.

Composition

Ingredients	Gms / Litre
MiVeg hydrolysate	20.00
Yeast extract	5.00
Sodium chloride	5.00
Sodium citrate	1.00
Esculin	1.00
Ferric ammonium citrate	0.50
Sodium azide	0.15
Final pH (at 25°C)	7.0 ± 0.2

** Formula adjusted, standardized to suit performance parameters.

Principle & Interpretation

Kanamycin Esculin Azide MiVeg Broth Base is prepared by adding MiVeg hydrolysate in place of Casein enzymic hydrolysate thus making the medium free from BSE/TSE risks. This medium is the modification of Kanamycin Esculin Azide Medium which was formulated by Mossel et al (1,2) for *Enterococci* detection in food stuffs. It can be used for bacteriological examination of food stuff using dip slide technique.

MiVeg hydrolysate and yeast extract supplies necessary nutrients for the growth of *Enterococci*. Kanamycin sulphate and sodium azide are the selective inhibitory components. Esculin and ferric ammonium citrate together form the indicator system to detect esculin-hydrolysing *Enterococci*.

Mossel et al described the following procedure as - 1gm/1ml food sample is added to prechilled diluent (Tryptone water, MiVeg VM1463) and decimal dilutions are prepared, which are then inoculated onto Kanamycin Esculin Azide MiVeg Broth Base (VM1776) and incubated at 35°C for upto 24 hours. Blackening of medium indicates esculin hydrolysis due to microbial action.

Methodology

Suspend 16.32 grams of powder media in 500 ml distilled water. Mix well and heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45–50°C and aseptically add rehydrated contents of one vial of Kanamycin Sulphate Selective Supplement (MS2146). Mix well before pouring into sterile tubes.

Warning: Sodium azide has a tendency to form explosive metal azides with plumbing materials thus it is advisable to use enough water to flush off the disposables.

Quality Control

Physical Appearance

Light yellow coloured, may have slightly greenish tinge, homogeneous, free flowing powder.

Colour and Clarity of prepared medium

Medium amber coloured, clear solution in tubes

Reaction

Reaction of 3.27% w/v aqueous solution is pH 7.0 ± 0.2 at 25°C.

pH Range

6.8 - 7.2

Cultural Response/Characteristics

Cultural characteristics observed after an incubation at 35°C or 42°C for 18-24 hours.

Organisms (ATCC)	Inoculum (CFU)	Growth*	Esculin hydrolysis
<i>Escherichia coli</i> (25922)	$>10^5$	inhibited	—
<i>Enterococcus faecium</i> (19434)	10^3 - 10^5	luxuriant	+
<i>Enterococcus bovis</i> (27960)	10^3 - 10^5	luxuriant	+
<i>Staphylococcus aureus</i> (25923)	10^3 - 10^5	inhibited	—
<i>Enterococcus faecalis</i> (29212)	10^3 - 10^5	luxuriant	+

KEY : * = with addition of Kanamycin Sulphate Selective Supplement (MS2146).

+ = Blackening of the medium / black zone around the colony.

- = No blackening of the medium / no black zone around the colony.

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 day.

Further Reading

1. Mossel D.A.A., Bijker, P.G.H. and Eelderink I., 1978, Arch. Lebensmittel - Hyg., 29:121.
2. Mossel D.A.A., et al., 1978, In : Streptococci., Skinner F.A. and Quesnel L. B. (ed.), SAB Symposium, series No.7, Academic Press, London.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.