

Technical Information

Reinforced Clostridial MiVeg Agar

Product Code :VM1154

Application:- Reinforced Clostridial MiVeg Agar is used for the cultivation and enumeration of *Clostridium* species and other anaerobes.

Composition

Ingredients	Gms / Litre
MiVeg hydrolysate	10.00
MiVeg extract	10.00
Yeast extract	3.00
Dextrose	5.00
Sodium chloride	5.00
Sodium acetate	3.00
Starch, soluble	1.00
L-Cysteine hydrochloride	0.50
Agar	13.50
Final pH (at 25°C)	6.8 ± 0.2

** Formula adjusted, standardized to suit performance parameters.

Principle & Interpretation

Reinforced Clostridial MiVeg Agar is prepared by adding MiVeg hydrolysate and MiVeg extract in place of Casein enzymic hydrolysate and beef extract thereby making the medium free from BSE/TSE risks. Reinforced Clostridial MiVeg Agar is the modification of Reinforced Clostridial Agar which was formulated by Hirsch and Grinstead (1). This media can be used to initiate growth from small inocula and to obtain the highest viable count of *Clostridia* and also in studies of spore forming anaerobes, especially *Clostridium butyricum* in cheese, for the enumeration of *Clostridia* in tube dilution counts and for preparation of plates for isolation (3). Other spore forming anaerobes, *Streptococci* and *Lactobacilli* can also be grown using this medium.

MiVeg hydrolysate, yeast extract, MiVeg extract and starch, supplies all the essential nutrients required for the optimum growth of *Clostridia*. Dextrose is the fermentable carbohydrate while sodium chloride maintains osmotic equilibrium of the medium. Cystine hydrochloride is the reducing agent whereas sodium acetate has buffering action. These media can be made selective by addition of 15-20mg Polymyxin B per litre of media (1).

Methodology

Suspend 51 grams of powder media in 1000 ml distilled water. Mix well and heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 10 lbs pressure (115°C) for 15 minutes.

Quality Control

Physical Appearance

Light yellow coloured, may have slightly greenish tinge, homogeneous, free flowing powder.

Gelling

Firm, comparable with 1.35% Agar gel.

Colour and Clarity of prepared medium

Light yellow coloured, clear to slightly opalescent gel forms in petri plates.

Reaction

Reaction of 5.1% w/v aqueous solution is pH 6.8 ± 0.2 at 25°C.

pH Range

6.6 - 7.0

Cultural Response/Characteristics

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours, in an anaerobic atmosphere.

Organisms (ATCC)	Inoculum (CFU)	Growth	Recovery
<i>Bacteroides fragilis</i> (23745)	10^2 - 10^3	good-luxuriant	>70%
<i>Bacteroides vulgatus</i> (8482)	10^2 - 10^3	good-luxuriant	>70%
<i>Clostridium butyricum</i> (9690)	10^2 - 10^3	good-luxuriant	>70%
<i>Clostridium perfringens</i> (13124)	10^2 - 10^3	good-luxuriant	>70%

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 day.



VM1154 Reinforced Clostridial MiVeg Agar

(Against dark background)

1. Control
2. *Bacteroides fragilis*

Further Reading

1. Hirsch and Grinsted, 1954, J. Dairy Res., 21:101.
2. Barnes and Ingram, 1956, J. Appl. Bact., 19:117.
3. Lewis and Angelotti (Eds.), 1964, Examination of Foods for Enteropathogenic and Indicator Bacteria, Dept. of HEW, PHS Publication, 1142, Washington.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.