

Technical Information

Halophilic MiVeg Broth

Product Code : VM1591

Application:- Halophilic MiVeg Broth is recommended for the isolation and cultivation of extreme halophiles.

Composition

Ingredients	Gms / Litre
MiVeg acid hydrolysate	10.00
Yeast extract	10.00
MiVeg peptone No.3	5.00
Trisodium citrate	3.00
Potassium chloride	2.00
Magnesium sulphate	25.00
Sodium chloride	250.00
Final pH (at 25°C)	7.2 ± 0.2

** Formula adjusted, standardized to suit performance parameters.

Principle & Interpretation

Halophilic MiVeg Broth is prepared by replacing animal based peptone with vegetable peptones thus making the medium free from BSE/TSE risks. Halophilic MiVeg Broth is the modification of Halophilic Broth which was formulated for isolation and cultivation of extreme halophilic species of *Halobacterium* and *Halococcus* from foods (1, 2). High salt concentration of about 20-30% is needed for optimum growth of these halophiles. These bacteria can cause pink discoloration on the outer surface accompanied by putrefaction and decomposition of fish, bacon and hides preserved in sea salts. MiVeg acid hydrolysate, MiVeg peptone No.3 and yeast extract supplies essential nutrients, mainly nitrogenous and vitamins for the optimum growth of the halophilic bacteria. Trisodium citrate serves as selective agent and also prevents loss of halophiles in mixed population; as it suppresses gram positive organisms and coliforms (2). Magnesium, an essential ion for the growth of extreme halophiles is incorporated in the medium as magnesium sulphate. 10 gm sample is added to 90 ml Halophilic MiVeg Broth and incubated at 35°C for upto 12 days.

Methodology

Suspend 30.5 grams of powder media in 100 ml distilled water. Mix thoroughly and heat to boiling to dissolve the medium completely. Dispense as desired and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Quality Control

Physical Appearance

Off white coloured, homogeneous, free flowing powder.

Colour and Clarity of prepared medium

Amber coloured, hazy solution containing heavy precipitate at the bottom in tubes.

Reaction

Reaction of 30.5% w/v aqueous solution is pH 7.2 ± 0.2 at 25°C.

pH Range

7.0 - 7.4

Cultural Response/Characteristics

Cultural characteristics observed after an incubation at 35-37°C for 12 days.

Organisms (ATCC)	Inoculum (CFU)	Growth	Recovery
<i>Halococcus morrhuae</i> (17082)	10^2 - 10^3	luxuriant	>70%
<i>Halobacterium salinarum</i> (33171)	10^2 - 10^3	luxuriant	>70%

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 day.

Further Reading

1. Dundas I.E., 1977, Advances In Microbiology and Physiology, Rose H. and Tempest D.W. (Eds.), A.P. London.
2. Gibbons N.E., 1969, Methods In Microbiology, Vol. 3B, Norris J.R., and Ribbons D.W. (Eds.), A.P., New York, pp.169-183.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.