

Technical Information

Bile Esculin MiVeg Agar Base

Product Code : VM1340

Application:- Bile Esculin MiVeg Agar Base is a differential medium recommended for isolation and presumptive identification of Group D *Streptococci* from food and pharmaceutical products.

Composition**

Ingredients	Gms / Litre
MiVeg peptone	22.00
MiVeg extract	6.00
Synthetic detergent No. II	5.00
MiVeg hydrolysate	15.00
Ferric citrate	0.50
Agar	15.00
Final pH (at 25°C)	6.6±0.2

** Formula adjusted, standardized to suit performance parameters.

Principle & Interpretation

Bile Esculin MiVeg Agar Base is prepared by using vegetable peptones instead of animal based peptones which makes the medium BSE/TSE risks free. This medium is the modification of Bile Esculin Agar which was formulated by Swan (1) for the isolation and identification of Group D *Streptococci* from food. Synthetic detergent No. II present in the medium inhibits gram positive bacteria other than group D *Streptococci* and *Enterococci*. *Enterococci* and Group D *Streptococci* were able to split esculin to esculetin and dextrose, which reacts with ferric citrate producing brownish black precipitate (2). Ferric citrate act as an indicator of esculin hydrolysis and resulting esculetin formation. Originally Bile Esculin Test was used for identification of *Enterococci*, but it was found that this test is also shared by Group D *Streptococci* (3) and therefore it is recommended that other tests such as salt tolerance be performed while identifying *Enterococci* (4). Similarly this medium was also shown to aid differentiation of *Enterobacteriaceae*, *Klebsiella*, *Enterobacter-Serratia* division from other *Enterobacteriaceae* genera (5) on the basis of esculin hydrolysis. Occasional strains of viridans *Streptococci* blacken the medium or display weakly positive reactions (6).

Methodology

Suspend 63.5 grams of powder media in 1000 ml purified/distilled water. Mix thoroughly. Heat to boiling to dissolve the medium completely. Add 1 gram of esculin (2 vials of MS2050). Mix & dispense into tubes or flasks as desired. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Allow the tubed medium to solidify in slanted position.

Quality Control

Physical Appearance

Brownish yellow coloured may have slightly greenish tinge, homogeneous, free flowing powder.

Gelling

Firm, comparable with 1.5% Agar gel.

Colour and Clarity of prepared medium

Yellow coloured, clear to slightly opalescent gel with a bluish tinge forms in petri plates.

Reaction

Reaction of 6.35 % w/v aqueous solution pH: 6.6±0.2 at 25°C

pH range

6.4-6.8

Cultural Response/Characteristics

Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours in an increased atmosphere of carbon dioxide

Organisms (ATCC)	Inoculum (CFU)	Growth	Recovery	Esculin hydrolysis
<i>Enterococcus faecalis</i> (29212)	10 ² -10 ³	luxuriant	>50%	+
<i>Streptococcus pyogenes</i> (19615)	10 ² -10 ³	None-poor	>10%	-
<i>Proteus mirabilis</i> (25933)	10 ² -10 ³	luxuriant	>50%	-


Key : + = Blackening of the medium

— = No Change

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.



VM1340 Bile Esculin MiVeg AgarBase

1. Control

2. *Enterococcus faecalis*

Further Reading

1. Swan A., 1954, J. Clin. Pathol., 7:160.
2. MacFaddin J.F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams and Wilkins, Baltimore.
3. Facklam R., 1972, Appl. Microbiol., 23:1131.
4. Facklam R., 1973, Appl. Microbiol., 26:138.
5. Edberg S.C., Pittman S., and Singer J.M., 1977, J. Clin. Microbiol., 6:111.
6. Facklam, et al 1999. In Murray, Baron, pfaller, Tenover and yolken (ed.), Manual of clinical Microbiology, 7th ed. ASM, Washington, D. C.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.