



Dehydrated Culture Media
Bases / Media Supplements

Technical Information

Pantothenate Assay Medium

Product Code: DM 1037

Application: - Pantothenate Assay Medium is recommended for the microbiological assay of Pantothenic acid or its salts using *Lactobacillus plantarum* ATCC 8014 as the test organism.

Composition**

Ingredients	Gms / Litre
Casein acid hydrolysate	10.000
Dextrose	40.000
Sodium acetate	20.000
L-Cystine	0.400
DL-Tryptophan	0.200
Adenine sulphate	0.020
Guanine hydrochloride	0.020
Uracil	0.020
Thiamine hydrochloride	0.0002
Riboflavin (Vitamin B2)	0.0004
Niacin	0.001
Pyridoxine	0.0008
p-Amino benzoic acid (PABA)	0.0002
Biotin	0.0000008
Monopotassium phosphate	1.000
Dipotassium phosphate	1.000
Magnesium sulphate	0.400
Sodium chloride	0.020
Ferrous sulphate	0.020
Manganese sulphate	0.020
Final pH (25°C)	6.8±0.2

**Formula adjusted, standardized to suit performance parameters



Principle & Interpretation

Pantothenate Assay Medium is prepared as per the recommendation of the U.S. Pharmacopeia ⁽¹⁾ for microbiological assay of pantothenic acid or its salts using *Lactobacillus plantarum* ATCC 8014 as the test organism.

This medium contains several nutrients and salts which are required for the growth of the test organism but not the pantothenic acid. Since the pantothenate is required for the growth of *Lactobacillus plantarum* assay strain, growth of the organism will occur only if the materials being assayed contain pantothenate. Exact concentration of pantothenate in the test material can be calculated by comparing results with standard curve of pantothenate.

Assay / Procedure: Stock cultures of the test organism: *Lactobacillus plantarum* ATCC 8014, are prepared in triplicate or more by stab inoculation of Pantothenate Culture Agar USP (DM1135). Following incubation for 16-24 hours at any selected temperature between 30°C and 37°C but held constant to within $\pm 0.5^\circ\text{C}$, the tubes are stored at 2-8°C. Prepare a fresh stab of stock culture every week as use a culture older than 1 week for transferring to broth for inoculation is not recommended for analyzing panto thenate.

Inoculum for the assay is prepared by subculturing from a suitable stock culture of *Lactobacillus plantarum* ATCC 8014 on Pantothenate Culture Agar USP (DM1135) into a tube containing 10 ml of sterile single strength Pantothenate Assay Medium supplemented with pantothenate. The medium is prepared by dissolving 36.5 grams of the dehydrated medium and 20 mcg of pantothenate in 1000 ml of distilled water. Distribute in tubes and sterilize by autoclaving for 15 minutes at 15 lbs pressure (121°C). After 18-24 hours incubation at 30-37°C the cells are centrifuged under aseptic conditions and the supernatant liquid is decanted. The cells are resuspended in 10 ml sterile 0.85% sodium chloride. The cell suspension is then diluted 1:100 with sterile 0.85% sodium chloride. The cell suspension so obtained is the inoculum. Inoculate each tube aseptically with 1 drop of the inoculum.

It is essential that a standard curve be set up for each assay since conditions of autoclaving, temperature of incubation, etc. which influence the standard curve readings, cannot be duplicated exactly from time to time. The standard curve is obtained by using calcium pantothenate solution at concentration of 0.0, 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09 and 0.1 mcg pantothenic acid per assay tube (10 ml). Turbidimetric determinations are made after 16-24 hours incubation at any selected temperature between 30°C and 37°C, but held constant to within $\pm 0.5^\circ\text{C}$. Acidimetric determinations are made after 72 hours incubation at 30-37°C. A standard curve is then plotted and concentration of unknown in the sample determined by extrapolation of standard curve.

Methodology

Suspend 7.31 grams of powder media in 100 ml distilled water. Shake well & heat to dissolve the medium completely and to distribute the slight precipitate evenly. Dispense in 5 ml amounts to each assay tube in increasing amounts of the standard or the unknown and total volume 10 ml per tube is adjusted by addition of distilled water. Sterilize by autoclaving at 15 lbs pressure (121°C) for 10 minutes. Cool the medium immediately. Generally satisfactory results are obtained when Calcium pantothenate at a concentration of 0, 0.025, 0.05, 0.075, 0.1, 0.125, 0.15 and 0.2 microgram per assay tube (10 ml) is used.

Quality Control

Physical Appearance

Off-white to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Light yellow coloured clear solution, which may have a slight precipitate.

Reaction

Reaction of 7.3% w/v aqueous solution at 25°C. pH : 6.8 \pm 0.2

pH range 6.60-7.00

Cultural Response/ characteristics

DM 1037: Microbiological Assay of Pantothenate is carried out by using *L. plantarum* ATCC 8014 after an incubation at 35-37°C for 18-24 hours.

Growth

Good growth is obtained. Gradually, increase in growth with increasing concentration of pantothenate standard level of 0.0, 0.025, 0.075, 0.1, 0.125, 0.15 and 0.2 mcg per assay tube is recorded as equivalent increase in absorbance at 620 nm.

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8°C in sealable plastic bags for 2-5 days.





Dehydrated Culture Media
Bases / Media Supplements

Further Reading

1. U.S. Pharmacopeia, National Formulary, 2002, 25/NF 20, U.S. Pharmacopoeial, Convention, Rockville, MD.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

