

Technical Information

Corn Meal Agar

Product Code: DM 1146

Application: - Corn Meal Agar is recommended for chlamyospore production by *Candida albicans* and the maintenance of fungal stock cultures.

Composition**

Ingredients	Gms / Litre
Corn meal, infusion from	50.000
Agar	15.000
Final pH (at 25°C)	6.0±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Corn Meal Agar is a well-established mycological medium used for the cultivation of fungi and to study chlamyospores production of *Candida* an accepted criterion for the identification of *Candida* species. Pollack and Benham ⁽¹⁾ have described the usefulness of this medium for studying the morphology of *Candida*. Walker and Huppert ⁽²⁾ modified this medium by adding polysorbate 80, which then stimulated formation of chlamyospore *Candida* species at a faster rate with increased number.

This has simple formulation containing only cornmeal infusion and agar which has enough nutrients to enhance the growth of fungi. Polysorbate 80 is a mixture of oleic esters, which activates the production of chlamyospore by *Candida albicans*, *Candida stellatoidea* and *Candida tropicalis* ⁽³⁾. Some *Candida* species lose their ability of chlamyospore formation by repeated sub culturing on this media

Pick a suspected colony from Sabouraud Dextrose Agar (DM1063) using a straight wire, and make a deep cut in the Corn Meal Agar plate. Repeat for each colony. Place a flamed sterile coverslip over the line of inoculum. After incubation for 24-48 hours at 25-30°C, the streaks are examined microscopically, through the coverslip, using low and high power objectives.

C. albicans produces mycelium bearing ball-like clusters of budding cells and characteristics thick walled round chlamyospores ⁽⁴⁾.

Methodology

Suspend 17 grams of powder media in 1000 ml distilled water. Shake well & heat to dissolve the medium completely. If desired add 1% polysorbate 80. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.

Quality Control

Physical Appearance

Cream to yellow coarse free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Light amber coloured, opalescent gel forms in Petri plates

Reaction

Reaction of 1.7% w/v aqueous solutions at 25°C. pH : 6.0±0.2

pH range 5.80-6.20

Cultural Response/ characteristics

DM 1146: Cultural characteristics observed after an incubation at 23-27°C for upto 4 days.



Dehydrated Culture Media
Bases / Media Supplements

Organism	Inoculum (CFU)	Growth	Chlamydo spores	Recovery
<i>Aspergillus brasiliensis</i> ATCC 16404	50-100	inhibited	negative	
<i>Candida albicans</i> ATCC 10231	50-100	inhibited	positive	>=70%
<i>Saccharomyces cerevisiae</i> ATCC 9763	50-100	luxuriant	negative	>=70%
<i>Saccharomyces uvarum</i> ATCC 28098	50-100	luxuriant	negative	>=70%

Key : * - Formerly known as *Aspergillus niger*

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Pollack and Benham, 1960, J. Lab. Clin. Med., 50:3 13.
2. Walker and Huppert, 1960, Tech. Bull. Reg. Med. Technol., 30:10.
3. Cooper and Silvo-Hunter, 1985, Manual of Clinical Microbiology, Lennette, Balows, Hausler and Shadomy (Eds.), 4th ed., ASM, Washington, D.C.
4. Conant N. F., Smith D. T., Baker R. D., Callaway J. L. and Martin D. S., 1971, Manual of Clinical Mycology, 3rd Ed., USA

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

