

Technical Information

Ayers & Johnson Agar (Stock Culture Agar)

Product Code: DM 1182

Application: Ayers and Johnson Agar (Stock Culture Agar) is recommended for the maintenance of cultures of Streptococci and other pathogenic and non pathogenic microorganisms.

Composition**

Ingredients	Gms / Litre
Beef heart, infusion from	500.000
Proteose peptone	10.000
Gelatin	10.000
Dextrose	0.500
Casein, purified	5.000
Disodium phosphate	4.000
Sodium citrate	3.000
Agar	7.500
Final pH (at 25°C)	7.5±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Maintenance medium are specially designed to maintain the viability of cultures for longer period. Stock Culture Agar was originally formulated by Ayers and Johnson ⁽¹⁾. They observed that in addition to supporting luxuriant growth, the medium also helped in maintaining the viability of Streptococci and various other organisms upto months together. They also observed that Streptococci maintained their viability as long as for four months when incubated in this medium at room temperature (25°C). Semisolid nature, a well-buffered environment and the presence of casein and dextrose, the latter, which serves as a source of energy helps the media in maintaining a viability of pathogenic organism Many fastidious organisms like *Mycobacterium* species, *S. pneumoniae*, show good growth on this medium. It can be made especially suitable for maintenance of Streptococci by the additions of L- Asparagine (1g/l) ⁽²⁾. Infusion from beef heart, proteose peptone, gelatin and casein serve as sources of nitrogen, vitamins and amino acids. Dextrose is a carbon and energy source. Disodium phosphate serves as a buffering agent while sodium citrate acts as a preservative.

Methodology

Suspend 50 grams of powder media in 1000 ml distilled water. Shake well and heat to boiling to dissolve the medium completely. Dispense in tubes. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Quality Control

Physical Appearance

Yellow to beige homogeneous coarse powder

Gelling

Semisolid, comparable with 0.75% Agar gel and 1.0% Gelatin gel.

Colour and Clarity of prepared medium

Light yellow coloured opalescent gel forms in tubes

Reaction

Reaction of 5% w/v aqueous solution at 25°C. pH : 7.5±0.2



Dehydrated Culture Media
Bases / Media Supplements

pH Range:- 7.30-7.70

Cultural Response/Characteristics

DM1182: Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours.

Organism	Inoculum (CFU)	Growth
<i>Neisseria meningitides</i> ATCC13090	50-100	Luxuriant
<i>Staphylococcus aureus</i> ATCC 25923	50-100	Luxuriant
<i>Staphylococcus pyogenes</i> ATCC 19615	50-100	luxuriant

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8⁰ in sealable plastic bags for 2-5 days.

Further Reading

1. Ayers and Johnson, 1924, J. Bacteriol., 9:111.
2. Atlas R. M., 2004, Handbook of Microbiological Media, 3rd Ed., CRC Press, Inc., Boca Raton, Fla.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.

