

Technical Information

Thioglycollate Medium w/o Dextrose

Product Code: DM 1190

Application: Thioglycollate Medium without Dextrose is used for cultivation of aerobes, microaerophiles, anaerobes and for fermentation studies with various carbohydrates.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	20.000
Sodium chloride	2.500
Dipotassium phosphate	1.500
Sodium thioglycollate	0.600
L-Cystine	0.400
Sodium sulphite	0.200
Methylene blue	0.002
Agar	0.500
Final pH (at 25°C)	7.2±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Thioglycollate Medium without Dextrose is the modification of original Thioglycollate medium ^(1, 2). This media used for the fermentation study of anaerobes and for enhancement of sporulation. Omission of dextrose from this media facilitates it to be used in fermentation studies with the addition of desired carbohydrate. Some Clostridia remain viable for a longer period and sporulate better in the absence of carbohydrate and thus this medium could be used for sporulations also.

Casein enzymic hydrolysate, L-cystine and salts provide essential nutrients like nitrogenous compounds, carbon, sulphur, minerals and amino acids. The reducing action provided by sodium thioglycollate and sodium sulphite binds molecular oxygen, thereby maintaining a low Eh ⁽³⁾. A small amount of agar is added to retard the absorption of oxygen by reducing convection currents in the medium. Methylene blue is a redox indicator.

Methodology

Suspend 25.7 grams of powder media in 1000 ml distilled water. If the medium is to be used for fermentation studies or for diagnostic work add 0.5 to 1% carbohydrate of choice. Shake well & heat to dissolve the medium completely. Dispense and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Alternatively, sterile carbohydrate solutions may be added to the broth after sterilization. The prepared medium should be stored in the dark at room temperature.

Note: If more than the upper one-third has acquired a green colour, the medium may be restored once by heating in a waterbath or free flowing steam until the green colour disappears.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Light yellow coloured very slightly opalescent viscous solution with upper 10% or less medium green on standing

Reaction

Reaction of 2.57% w/v aqueous solution at 25°C. pH : 7.2±0.2



Dehydrated Culture Media
Bases / Media Supplements

pH Range:- 7.00-7.40

Cultural Response/Characteristics

DM 1190: Cultural characteristics observed after an incubation at 35-37°C for 48 hours (in an appropriate atmosphere).

Organism	Inoculum (CFU)	Growth
<i>Bacillus subtilis</i> ATCC 6633	50-100	good
<i>Bacteroides vulgatus</i> ATCC 8482	50-100	fair
<i>Candida albi cans</i> ATCC 10231	50-100	good
<i>Clostridium sporogenes</i> ATCC 11437	50-100	good-luxuriant
<i>Micrococcus luteus</i> ATCC 10240	50-100	good
<i>Neisseria meningitidis</i> ATCC 13090	50-100	good
<i>Streptococcus pyogenes</i> ATCC 19615	50-100	good-luxuriant

Storage and Shelf Life-

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8⁰ in sealable plastic bags for 2-5 days.

Further Reading

1. Brewer J. H., 1940, J. Am Med. Assoc., 115, 598.
2. Brewer J. H., 1940, J. Bacteriol., 39:10.
3. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol.1 William and Wilkins, Baltimore.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performs parameters.

