

Technical Information

Brain Heart Infusion Broth

Product Code: DM 1210I

Application: Brain Heat Infusion Broth is highly nutritious and is recommended for the propagation of pathogenic cocci and other fastidious organisms associated with blood culture work and allied pathological investigations and for enrichment of *Staphylococcus aureus*.

Composition**

Ingredients	Gms / Litre
Peptic digest of animal tissue	10.000
Calf brain, infusion (solids)	12.500
Beef heart, infusion (solids)	5.000
Dextrose	2.000
Sodium chloride	5.000
Disodium phosphate	2.500
Final pH (at 25°C)	7.4±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Rosenow ⁽¹⁾ devised the original Brain Heart Infusion Broth by adding brain tissue to dextrose broth. Brain Heart Infusion Broth is a highly nutritious medium and is also well buffered to support the growth of a number of microorganisms ⁽²⁻⁴⁾. Recently this medium has been recommended by ISO committee for the detection of *Staphylococcus aureus* ⁽⁵⁾. With the additions of desired additives this medium can be specifically adopted for cultivation of different bacteria. Addition of 6.5% sodium chloride makes it selective for isolation of various salt tolerant bacteria like Enterococci. Brain Heart Infusion Broth is also used for the preparation of inocula in antimicrobial susceptibility tests.

Methodology

Suspend 37 grams of powder media in 1000 ml distilled water. Dispense into bottles or tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. For best results, the medium should be used on the day it is prepared, otherwise, it should be boiled or steamed for a few minutes and then cooled before use.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Light amber coloured, clear to slightly opalescent solution

Reaction

Reaction of 3.7% w/v aqueous solution at 25°C. pH : 7.4±0.2

pH range 7.20-7.60

Cultural Response/Characteristics

DM1210I: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.



Dehydrated Culture Media
Bases / Media Supplements

Organism	Inoculum (CFU)	Growth
<i>Neisseria meningitides</i> ATCC 13090	50-100	Luxuriant
<i>Staphylococcus aureus</i> ATCC 25923	50-100	Luxuriant
<i>Streptococcus pneumonia</i> ATCC 6303	50-100	Luxuriant
<i>Streptococcus pyogenes</i> ATCC 19615	50-100	Luxuriant

Storage and Shelf Life

Dried media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8^o in sealable plastic bags for 2-5 days.

Further Reading

1. Rosenow, 1919, J. Dental Res., 1:205.
2. Roseburg T. et al, 1944, J. Inf. Dis., 74:13 1.
3. Conant N.F., 1950, Diagnostic Procedures and Reagents, 3rd Ed., APHA, Inc., New York, p. 452.
4. MacFaddin J.F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.
5. International Organization for Standardization (ISO), 1983, Enumeration of Staphylococcus aureus, Draft ISO/DIS 6880.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

