

Technical Information

Deoxycholate Citrate Agar without Sucrose

Product Code: DM1222

Application: Deoxycholate Citrate Agar without Sucrose is used for differentiation and identification of enteric pathogens.

Composition**

Ingredients	Gms / Litre
Biopeptone	7.000
Meat extract	3.000
Sodium deoxycholate	2.500
Sodium citrate	10.500
Lactose	5.000
Sodium thiosulphate	5.000
Neutral red	0.030
Agar	12.000
Final pH (at 25°C)	7.2±0.2

**Formula adjusted, standardized to suit performance parameters.

Principle & Interpretation

Deoxycholate Citrate Agar without Sucrose is used for differentiation and identification of members of family *Enterobacteriaceae*. Leifson ⁽¹⁾ developed Deoxycholate Agar as a differential medium containing pure chemicals. Deoxycholate Citrate Agar without Sucrose contains biopeptone and meat extract, which supply essential nutrients for the support of bacterial growth. Citrate and deoxycholate act as inhibitors. Sodium deoxycholate and sodium citrate inhibit gram-positive organisms. Lactose helps in differentiating enteric bacilli as lactose fermenters produce red coloured colonies while lactose non-fermenters form colourless colonies. Citrate and iron (Fe) combination has a strong hydrolyzing effect on agar when the medium is heated, producing a soft and unelastic agar. If autoclaved the agar becomes soft and almost impossible to streak ⁽¹⁾.

Methodology

Suspend 45.03 grams of powder media in 1000 ml distilled water. Shake well and heat, to dissolve the medium completely. DO NOT OVERHEAT OR AUTOCLAVE.

Quality Control

Physical Appearance

Light yellow to pink homogeneous free flowing powder

Gelling

Firm, comparable with 1.2% Agar gel.

Colour and Clarity of prepared medium

Reddish orange coloured, clear to slightly opalescent gel forms in Petri plates.

Reaction

Reaction of 4.5% w/v aqueous solution at 25°C. pH : 7.2±0.2

pH Range 7.00-7.40

Cultural Response/Characteristics

DM1222: Cultural characteristics observed after an incubation at 35 - 37°C for 18 - 24 hours.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of Colony
<i>Bacillus subtilis</i> ATCC 6633	$\geq 10^3$	Inhibited	0%	
<i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant	$\geq 50\%$	pink with bile precipitate
<i>Enterobacter aerogenes</i> ATCC 13048	50-100	good-luxuriant	$\geq 50\%$	Pink
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	good-luxuriant	$\geq 50\%$	colourless
<i>Enterococcus faecalis</i> ATCC 29212	$\geq 10^3$	Inhibited	0%	

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8^o in sealable plastic bags for 2-5 days.

Further Reading

1. Leifson, 1935 J. Path. Bacteriol, 40:58 1.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.