

## Technical Information

### Motility Test Medium

**Product Code: DM1260**

**Application:** - Motility Test Medium is recommended for detection of bacterial motility.

### Composition\*\*

Ingredients	Gms / Litre
Tryptose	10.000
Sodium chloride	5.000
Agar	5.000
Final pH (25°C)	00207.2±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Principle & Interpretation

Bacterial motility can be observed directly on microscopic slide or it can be seen on motility media having agar concentration of 0.4% or less <sup>(1)</sup>. Use of such semisolid media to observe or detect motility was reported by Tittsler and Sandholzer <sup>(2)</sup>. Motility Test Medium is a modification of their formulation in which Motility can be visualized as a diffused zone of growth flaring out from the line of inoculation <sup>(1)</sup>. Hanging-drop technique in motility tests has practical difficulties, which is efficiently eliminated by use of culture-based methods using semi-solid media.

Tryptose serve as a source of essential growth nutrients required for bacterial metabolism. Sodium chloride maintains the osmotic equilibrium of the medium. Small amount of agar helps to create a semisolid medium.

Bacterial motility can be observed directly by examination of the tubes following incubation. Inoculation is done by stabbing through the centre of the medium. Incubate at appropriate temperature for 18-40 hours. Non-motile organisms grow only along the line of inoculation whereas motile organisms grow away from the line of inoculation or may show growth even throughout the medium. Weak or equivocal motility results should be confirmed by flagellum stain or by direct wet microscopy (hanging drop) <sup>(3, 4)</sup>.

### Methodology

Suspend 20 grams of powder media in 1000 ml distilled water. Shake well & heat to dissolve the medium completely. Dispense in tubes and sterilize by autoclaving at 15 lbs pressure (12 1°C) for 15 minutes. Allow tubed medium to cool in an upright position.

### Quality Control

#### Physical Appearance

Cream to yellow homogeneous free flowing powder

#### Gelling

Semisolid, comparable with 0.5% Agar gel.

#### Colour and Clarity of prepared medium

Light yellow coloured clear to slightly opalescent gel forms in tubes as butts

#### Reaction

Reaction of 2.0% w/v aqueous solution at 25°C. pH : 7.2±0.2

**pH range** 7.00-7.40

#### Cultural Response/ characteristics

DM 1260: Cultural characteristics observed after an incubation at 35 - 37°C for 18 - 48 hours.



Dehydrated Culture Media  
Bases / Media Supplements

Organism	Inoculum (CFU)	Growth	Motility
Escherichia coli ATCC 25922	50-100	luxuriant	positive, growth away from stabline causing turbidity
Enterobacter aerogenes ATCC 13048	50-100	luxuriant	positive, growth away from stabline causing turbidity
Klebsiella pneumoniae ATCC 13883	50-100	luxuriant	negative, growth the stabline, surrounding medium remains clear
Salmonella Enteritidis ATCC 13076	50-100	luxuriant	positive, growth away from stabline causing turbidity
Staphylococcus aureus ATCC 25923	50-100	luxuriant	negative, growth the stabline, surrounding medium remains clear

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

## Further Reading

1. Koneman E. W., Allen S. D., Janda W. M., Schreckenberger P. C., Winn W. C. Jr., (Eds.), 1992, Colour Atlas and Textbook of Diagnostic Microbiology, 4th Ed., J. B. Lippincott Company.
2. Tittsler R. P. and Sandholzer L. A., 1936, J. Bacteriol., 3 1:575.
3. DAmato R. F., and Tomfohrde K. M., 1981, J. Clin. Microbiol., 14 (3), 347-348.
4. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams and Wilkins, Baltimore

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

