

Technical Information

Phenol Red Lactose Agar

Product Code: DM 1270

Application: Nutrient Agar, pH 6.0 with 0.8% NaCl is used as a general purpose culture media.

Composition**

Ingredients	Gms / Litre	
Proteose peptone	10.000	
Beef extract	1.000	
Sodium chloride	5.000	
Lactose	10.000	
Phenol red	0.025	
Agar	15.000	
Final pH (at 25°C)	7.4±0.2	

^{**}Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Phenol Red Agar media are recommended ⁽¹⁻³⁾ for studying the fermentation of various carbohydrates individually using the pure cultures.

Proteose peptone which is free from fermentable carbohydrates is added in the medium thereby preventing the occurance of false positive reactions. When Phenol Red Agar with Lactose is used, a positive carbohydrate fermentation reaction is indicated by the production of a yellow colour in agar due to production of acid. Gas production is indicated by the splitting of agar or by the bubbles formation. Plates or tubes may be incubated aerobically or anaerobically depending on the type of the test organism.

Methodology

Suspend 41.02 grams of powder media in 1000 ml distilled water. Shake well & heat with frequent agitation to dissolve the medium completely. Dispense in tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Allow the tubed media to cool in slanted position to form slants with deep butts.

Quality Control

Physical Appearance

Light yellow to pink homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Red coloured clear to slightly opalescent gel forms in tubes as slants

Reaction

Reaction of 4.1% w/v aqueous solution at 25°C. pH: 7.4±0.2 pH: 7.4±0.2

pH Range 7.20-7.60

Cultural Response/ characteristices

DM 1270: Cultural characteristics observed after an incubation at 35 37°C for 18 - 24 hours

Organism	Inoculum (CFU)	Growth	Acid	Gas
Alcaligenes faecalis ATCC 8750	50-100	luxuriant	Negative reaction, no colour chan	ge Negative reaction
Escherichia coli ATCC 25922	50-100	luxuriant	Positive reaction, yellow colour	Positive reaction
Klebsiella pneumoniae ATCC 13883	50-100	luxuriant	Positive reaction, yellow colour	Positive reaction





Proteus vulgaris ATCC 13315	50-100	luxuriant	Negative reaction, no colour change Negative reaction
Salmonella Typhimurium ATCC 14028	50-100	luxuriant	Negative reaction, no colour change Negative reaction
Shigella flexneri ATCC 12022	50-100	luxuriant	Negative reaction, no colour change Negative reactio

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label. **Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

Further Reading

- 1. MacFaddin J., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins,
- 2. Finegold and Baron, 1986, Bailey and Scotts Diagnostic Microbiology, 7th ed., The C.V. Mosby Co., St. Louis.
- 3. Ewing, 1986, Edwards and Ewings Identification of Enterobacteriaceae, 4th ed., Elsevier Science Publishing Co., Inc., New York.

Disclaimer:

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performens parameters.

