

Technical Information

Sulphate API Broth w/o Sodium Lactate

Product Code: DM 1310

Application: Sulphate API Broth is used for detection and estimation of sulphate reducing bacteria.

Composition**

Ingredients	Gms / Litre
Yeast extract	1.000
Magnesium sulphate	0.200
Dipotassium phosphate	0.010
Ferrous ammonium sulphate	0.100
Sodium chloride	10.000
Final pH (at 25°C)	7.5±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Sulphate API Broth is prepared according to the formulation of the American Petroleum Institute Recommended Practice ⁽¹⁾ for detection of sulphate reducing bacteria. Sulphate-reducing bacteria cause corrosion of oil well systems resulting in perforations in the pipes. Sulphate-reducing bacteria convert sulphate to sulphide which with the ferrous ion gives black colour. The insoluble sulphide results in plugging of pipe.

Yeast extract in the medium provides nitrogen and other nutrients necessary to support bacterial growth. Ascorbic acid is the carbohydrate source. Dipotassium Phosphate buffers the medium. Sodium chloride, magnesium sulphate and ferrous ammonium sulphate provide essential ions. *Desulfovibrio* oxidizes reduced substrates i.e. sodium lactate, further with stepwise reduction of sulfate to sulfide. The detection and estimation of these bacteria is done on the basis of their ability to grow and produce sulphide in this medium. For the estimation, appropriate dilutions of water samples are inoculated in the test proper.

Methodology

Suspend 11.41 grams of powder media in 1000 ml distilled water. Add 4 ml of sodium lactate. Shake well & heat if necessary to dissolve the medium completely. Dispense, preferably in screw-capped tubes in 9 ml amounts. Sterilize by autoclaving at 15 lbs pressure (121°C) for 10 minutes. Close the caps immediately while the medium is still hot.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Light yellow coloured clear solution in tubes

Reaction

Reaction of 1.1% w/v aqueous solutions (containing 0.4% v/v Sodium lactate) at 25°C. pH : 7.5±0.2

pH range: 7.30-7.70

Cultural Response/Characteristics

DM 1310: Cultural characteristics observed after an incubation at 30°C for upto 1 week, under anaerobic condition.

Organism

Inoculum (CFU)

Growth

Desulfovibrio desulfuri cans ATCC 13541

50-100

good-luxuriant



Dehydrated Culture Media
Bases / Media Supplements

Storage and Shelf Life

Dried media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8⁰ in sealable plastic bags for 2-5 days.

Further Reading

1. American Petroleum Institute Recommended Practice 28, 1959, First ed.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

