

## Technical Information

### Bushnell Haas Agar

#### Product Code: DM 1349

**Application:** - Bushnell Haas Agar is recommended for the microbial examination of fuels and for studying microbial hydrocarbon deterioration.

#### Composition\*\*

| Ingredients             | Gms / Litre |
|-------------------------|-------------|
| Magnesium sulphate      | 0.200       |
| Calcium chloride        | 0.020       |
| Monopotassium phosphate | 1.000       |
| Dipotassium phosphate   | 1.000       |
| Ammonium nitrate        | 1.000       |
| Ferric chloride         | 0.050       |
| Agar                    | 20.000      |
| Final pH ( at 25°C)     | 7.0±0.2     |

\*\*Formula adjusted, standardized to suit performance parameters

#### Principle & Interpretation

Bushnell Haas Agar is prepared according to the formula of Bushnell and Haas <sup>(1)</sup> and recommended for the microbiological examination of fuels by the SIM Committee on microbiological deteriorations of fuels <sup>(2)</sup>. These media contain all nutrients except carbon source, necessary for the growth of bacteria. Only those bacteria that are able to decompose hydrocarbon will grow in these media. Specific carbon source i.e. (hydrocarbon) can be added to this medium and their utilization by different microorganisms can be studied.

These bacteria can decompose hydrocarbons like kerosene, mineral oil, paraffin wax and gasoline. Liquid hydrocarbon is layered on the surface of inoculated agar. For testing volatile hydrocarbons such as gasoline the Petri-plates containing the medium are inverted and the hydrocarbon is poured into the lid. Magnesium sulphate, calcium chloride and ferric chloride provide trace elements. Ammonium nitrate is a nitrogen source while monopotassium phosphate and potassium phosphate buffers the medium.

#### Methodology

Suspend 23.27 grams of powder in 1000 ml distilled water. Shake well & heat to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. A white precipitate prior to sterilization becoming yellow to orange after sterilization is normal.

#### Quality Control

##### Physical Appearance

White to cream homogeneous free flowing powder

##### Gelling

Firm, comparable with 2.0% agar gel.

##### Colour and Clarity of prepared medium

Light amber coloured, clear to slightly opalescent gel forms in Petri plates.



Dehydrated Culture Media  
Bases / Media Supplements

#### Reaction

Reaction of 2.33% w/v aqueous solution at 25°C. pH : 7.0±0.2

**pH range** 6.80-7.20

#### Cultural Response/ characteristics

**DM 1349:** Cultural characteristics observed after an incubation at 25-30°C within 1 week.

| Organism                          | Inoculum (CFU) | Growth (Plain) | Growth w/ minerals |
|-----------------------------------|----------------|----------------|--------------------|
| Pseudomonas aeruginosa ATCC 27853 | 50-100         | poor           | good-luxuriant     |
| Pseudomonas aeruginosa A TCC 9027 | 50-100         | poor           | good-luxuriant     |

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

## Further Reading

1. Bushnell and Haas, 1941, J. Bacteriol., 41:653.
2. Allred, DeGray, Edwards, Hedrick, Klemme, Rogers, Wulf and Hodge, 1963, Proposed Procedures for Microbiological Examination of Fuels, SIM Special Publications, No. 1. Merck, Sharp & Dohme Research Laboratories, Rahway, N.J.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

