

Technical Information

Peptone Iron Agar

Product Code: DM 1440

Application: Peptone Iron Agar is used for detection of hydrogen sulfide production by microorganisms.

Composition**

Ingredients	Gms / Litre
Peptic digest of animal tissue	15.000
Proteose peptone	5.000
Ferric ammonium citrate	0.500
Sodium glycerophosphate	1.000
Sodium thiosulphate	0.080
Agar	15.000
Final pH (at 25°C)	6.7±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

The ability of certain bacterial species to liberate sulfur from sulfur-containing amino acids or other compounds in the form of hydrogen sulphide is an important characteristic for their identification. Hydrogen sulphide production can be detected by adding a sulfur source and an H₂S indicator system in the medium ⁽¹⁾. Peptone Iron Agar which is modification of Levin's original formula ^(2, 3) is used to detect H₂S production by organisms. This medium utilizes sodium thiosulphate, an inorganic compound as a supplemental source of sulfur and ferric ammonium citrate as the H₂S indicator in the medium. Peptone Iron Agar is more accurate over Lead Acetate Agar, medium for detecting and giving clear and early results for H₂S producing organism. ⁽⁴⁾. This is because ferric ammonium citrate is a better indicator of hydrogen sulphide, as compared to lead acetate.

Peptic digest of animal tissue and proteose peptone provide carbonaceous and nitrogenous compounds, including trace elements. Sodium thiosulphate and ferric ammonium citrate form the H₂S detecting system. Sulphide is released from thiosulphate due to the action of bacterial enzymes. This sulphide then couples with a hydrogen ion to form H₂S, which then reacts with the ferric ions from ferric ammonium citrate to produce insoluble heavy metal sulphides that appear as a black precipitate ⁽¹⁾. Sodium glycerophosphate buffers the medium.

Methodology

Suspend 36.58 grams of powder media in 1000 ml distilled water. Shake well & heat to dissolve the medium completely. Dispense in test tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Allow the tubed medium to cool in an upright position or in a slanting position to form slants.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Light amber coloured clear to slightly opalescent gel forms in tubes as slants



Dehydrated Culture Media
Bases / Media Supplements

Reaction

Reaction of 3.66% w/v aqueous solution at 25°C. pH : 6.7±0.2

pH range 6.50-6.90

Cultural Response/Characteristics

DM 1440: Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours.

Organism	Inoculum (CFU)	Growth
<i>Escherichia coli</i> ATCC 25922	50-100	negative reaction, no blackening of medium
<i>Enterobacter aerogenes</i> ATCC 13048	50-100	negative reaction, no blackening of medium
<i>Salmonella Typhi</i> ATCC 6539	50-100	positive reaction, blackening of medium
<i>Salmonella Enteritidis</i> ATCC 13076	50-100	positive reaction, blackening of medium

Storage and Shelf Life

Dried media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Koneman E. W, Allen S. D., Janda W. M., Schreckenberger P. C., Winn W.C. Jr., 1992, Colour Atlas and Textbook of Diagnostic Microbiology, 4th Ed, J. B. Lippincott Company, Philadelphia.
2. Levine M., Vaughn R., Epstein S. S. and Anderson D., 1932, Proc. Soc. Exp. Biol. Med. 29:1022.
3. Levine M., Epstein S. S. and Vaughn R., 1934, Am. J. Public Health 24 :505.
4. Tittsler R. P. and Sandholzer L. A., 1937, Am. J. Public Health 27: 1240.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

