

Technical Information

Motility Medium S Base

Product Code: DM1514

Application: - Motility Medium S Base is recommended for easy detection of bacterial motility by means of TTC reduction.

Composition**

Ingredients	Gms / Litre
Beef heart, infusion from	500.000
Tryptose	10.000
Gelatin	30.000
Sodium chloride	5.000
Dipotassium phosphate	2.000
Potassium nitrate	2.000
Agar	1.000
Final pH (25°C)	7.3±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Bacterial motility is an important feature in making a final species identification. Bacteria move by means of flagella, the number and location of which may vary among different species. Interpretation of motility test is done by macroscopic examination of the motility medium for a diffused zone of growth flaring out from the line of inoculation ⁽¹⁾.

Motility Medium S Base formulated as per Ball and Sellers ⁽²⁾, a multipurpose media is used to determine motility, gelatin liquefaction and nitrate reduction. All the tests can be determined using a single tube.

Beef heart infusion and tryptose provide nitrogenous compounds, sulphur, carbon and other essential growth nutrients. Sodium chloride maintains osmotic equilibrium. 0.1% agar in presence of 3% gelatin is sufficient to preserve an intact stab line. Nitrate reduction is tested using nitrate reagents after recording motility results ⁽³⁾. Motility is observed as diffused growth away from the stab inoculation line while non-motile organisms grow along the stab line. The use of TTC aids in the visual detection of bacterial motility. Tetrazolium salts are colourless but are converted into insoluble red formazan complexes by the reducing properties of growing bacteria. Development of this red colour helps to trace the spread of bacteria from the inoculation line. However these salts may inhibit certain fastidious bacteria and cannot be used in all cases ⁽¹⁾. Potassium nitrate serves as a substrate for nitrate reaction. Organisms capable of reducing nitrate exhibit increased motility in presence of 0.2% potassium nitrate, especially nitrate-reducing obligate aerobes ⁽²⁾. Phosphate maintains buffering in the medium and it also has a stimulatory effect on motility of Proteus species ⁽²⁾. Organisms having the ability to produce gelatinase digest or liquefy gelatin. Gelatin liquefaction can be determined by placing the test medium tubes in a refrigerator or ice bath, after an incubation at 35-37°C ⁽⁴⁾.

Methodology

Suspend 60 grams of powder media in 1000 ml distilled water. Shake well and heat to boil to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45 - 50°C and add 10 ml of 1% solution of 2, 3, 5-Triphenyl Tetrazolium Chloride (TTC Solution 1%, MS2057). Mix well and dispense in sterile tubes.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Semisolid, comparable with 0.1% Agar gel and 3.0% Gelatin gel.

Colour and Clarity of prepared medium

Yellow coloured clear to slightly opalescent gel forms in tubes as butts

Reaction

Reaction of 6.0% w/v aqueous solution at 25°C. pH: 7.3±0.2

pH range 7.10-7.50

Cultural Response/ characteristics

DM 1514: Cultural characteristics observed with added 1% TTC solution (MS2057) after an incubation at 35-37°C for 18-48 hours.

Organism	Inoculum (CFU)	Growth	Motility	TTC reduction
<i>Enterobacter aerogenes</i> ATCC 13048	50-100	good-luxuriant	positive growth away from stabline maroon turbidity	positive reaction, red to maroon colour
<i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant	positive growth away from stabline maroon turbidity	positive reaction, red to maroon colour
<i>Klebsiella pneumoniae</i> ATCC 13883	50-100	good-luxuriant	negative, growth along reaction, the stabline, surrounding medium remains clear.	positive reaction, red to maroon colour
<i>Proteus mirabilis</i> ATCC 25933	50-100	good-luxuriant	negative, motility is temperature dependent. It is more pronounced at 20°C and almost absent at 35°C	positive reaction, red to maroon colour

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Koneman E. W., Allen S. D., Janda W. M., Schreckenberger P. C., Winn W. C. Jr., 1992, Colour Atlas and Textbook of Diagnostic Microbiology, 4th Ed., J. B. Lippincott Company
2. Ball R. J. and Sellar W., 1966, Appl. Microbiol., 14 (4): 670.
3. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.
4. MacFaddin J. F., 2000, Biochemical tests for Identification of Medical Bacteria, 3rd Ed., Lippincott, Williams and Wilkins, Baltimore.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.