

## Technical Information

### MacConkey Broth (Double strength) w/ Neutral Red

#### Product Code: DM 1539

**Application:** - MacConkey Broth (Double strength) w/ Neutral Red is recommended for the primary isolation of coliforms from large samples such as water and wastewater.

#### Composition\*\*

Ingredients	Gms / Litre
Peptic digest of animal tissue	40.000
Lactose	20.000
Bile salts	10.000
Sodium chloride	10.000
Neutral red	0.150
Final pH (at 25°C)	7.4±0.2

\*\*Formula adjusted, standardized to suit performance parameters

#### Principle & Interpretation

MacConkey Broth is widely used as a differential medium for isolation detection and enumeration of coliforms from a wide variety of clinical, food and water samples. Identification of coliform is based on colour change of the medium due to the presence of the indicator neutral red <sup>(1, 2)</sup>. Peptic digest of animal tissue provides necessary nitrogen source. Lactose serves as the fermentable carbohydrate source. Sodium chloride maintains the osmotic balance of the cells. The selective action of these media is due to the presence of bile salts, which are inhibitory to most species of gram-positive bacteria. Gram-negative bacteria usually grow well on these media and are differentiated by their ability to ferment lactose. The colour change of the medium shown by lactose-fermenters is due to production of acid from lactose and a subsequent colour change of the indicator dye when the pH of the media falls below 6.8. Lactose non-fermenting strains, such as *Shigella* and *Salmonella* do not alter the appearance of the media.

MacConkey Broth (Double Strength) w/ Neutral Red DM1539 is recommended for the primary isolation of coliforms from large samples such as water and wastewater. The medium turns pink in case of lactose fermentors and yellow in case of lactose- nonfermenters, due to neutral red. MacConkey Broth Double Strength w/Neutral Red DM1539 has the same composition in double strength to that of MacConkey Broth (DM1007), which contains neutral red as an indicator and is considered as a standard medium for the primary isolation as well as presumptive identification of coliform-aerogenes group of organisms in food and water.

#### Methodology

Suspend 80.15 grams of powder media in 1000 ml distilled water. Shake well & heat if necessary to dissolve the medium completely. Distribute into test tubes with inverted Durham tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool the tubes before inoculation.

#### Quality Control

##### Physical Appearance

Light yellow to pink homogeneous free flowing powder

##### Colour and Clarity of prepared medium

Red coloured clear solution without any precipitate

##### Reaction

Reaction of 8.02% w/v aqueous solution at 25°C. pH : 7.4±0.2

pH range 7.20-7.60

## Cultural Response/characteristics

**DM 1539:** Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth	Acid	Gas
Escherichia coli ATCC 25922	50-100	luxuriant	positive reaction	positive reaction
Enterobacter aerogenes ATCC 13048	50-100	luxuriant	positive reaction	positive reaction
Salmonella Choleraesuis ATCC 12011	50-100	fair - good	negative reaction	negative reaction
Staphylococcus aureus ATCC 25923	$\geq 10^3$	inhibited		

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

## Further Reading

1. MacConkey, 1900, The Lancet, ii:20.
2. MacConkey, 1905, J. Hyg., 5:333.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.