



Dehydrated Culture Media
Bases / Media Supplements

Technical Information

Pantothenate Inoculum Broth

Product Code: DM 1542

Application: Pantothenate Inoculum Broth is recommended for the preparation of inoculum used in microbiological assays of pantothenic acid or its salts.

Composition**

Ingredients	Gms / Litre
Peptonized milk	15.000
Yeast extract	5.000
Dextrose	10.000
Monopotassium phosphate	2.000
Tomato juice (100 ml)	5.000
Polysorbate 80	1.000
Final pH (at 25°C)	6.8±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Pantothenate Inoculum Broth is prepared based on the formula originally designed by Kulp and White ⁽¹⁾ and later on modified and recommended by AOAC ⁽²⁾ to obtain high recovery of *Lactobacillus acidophilus* used in microbiological assays. Kulp and White developed this medium for plating ⁽¹⁾ of *Lactobacillus acidophilus*.

Peptonized milk contains lactose, which serves as the energy source for *Lactobacilli*. Dextrose serves as the fermentable carbohydrate and/or energy source. Yeast extract and peptone (of peptonized milk) provide vitamin B complex, nitrogenous compounds and trace ingredients for the growth. Tomato juice provides an acid environment in the medium resulting in inhibition of microorganisms other than acidophilic bacteria. Polysorbate 80 supplies fatty acids required for the metabolism of *Lactobacilli*.

Methodology

Suspend 38 grams of powder media in 1000 ml distilled water. Shake well and heat if necessary to dissolve the medium completely. Distribute in tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Medium amber coloured clear solution, without any precipitate.

Reaction

Reaction of 3.8% w/v aqueous solution at 25°C. pH : 6.8±0.2

pH range: 6.6-7.0

Cultural Response/ characteristics

DM 1542: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth
<i>Lactobacillus casei</i> ATCC 9595	50-100	luxuriant
<i>Lactobacillus leichmannii</i> ATCC 4797	50-100	luxuriant
<i>Lactobacillus plantarum</i> ATCC 8014	50-100	luxuriant
<i>Lactobacillus acidophilus</i> ATCC 11506	50-100	luxuriant





Dehydrated Culture Media
Bases / Media Supplements

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Kulp J. W. L. and White V., 1932, Science, 76:17.
2. Cunniff P., (Ed.), 1995, Official Methods of Analysis of AOAC International, 16th Ed., Vol. II, AOAC, Arlington, Virginia.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

