

Technical Information

Buffered Yeast Agar

Product Code: DM 1585

Application: - Buffered Yeast Agar is used (I) as a semisynthetic medium for the cultivation of yeasts and moulds and (II) for controlling bottle washing operations in soft drinks and related industries.

Composition**

Ingredients	Gms / Litre
Yeast extract	5.000
Dextrose	20.000
Ammonium sulphate	0.720
Ammonium dihydrogen phosphate	0.260
Agar	15.000
Final pH (at 25°C)	5.5±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Yeasts grow well on a minimal medium containing only dextrose and salts. The addition of yeast extract allows faster growth so that during exponential or log phase growth, the cells divide every 90 minutes ⁽¹⁾. Buffered Yeast Agar is prepared as per the modification of the yeast-salt medium as described by Davis else where ⁽²⁾.

The medium contains yeast extract, which supplies B-complex vitamins to stimulate growth. Dextrose is the carbohydrate source. The reaction of this medium can be adjusted to required pH values by the addition of citric or lactic acid to the medium after sterilization. The following table shows the amount of the acids required to be added to 100 ml of Buffered Yeast Agar cooled to 50°C.

Volume of acid to be added to 100 ml of medium to achieve the desired pH

pH	1% w/v solution of Citric acid monohydrate (ml)	1% w/v solution of Lactic acid (ml)
4.75	1.26	0.125
4.5	2.24	0.2
4.25	3.92	0.3
4.0	6.16	0.45
3.75	9.52	0.7
3.5	14.56	1.17

Bunker ^(3, 4) described a practical method for assessing the efficiency of the bottle cleaning operations. In this method, the bottle under test is converted into a roll-tube culture by coating tubes or botals internally with the medium. When the agar sets, the tube/bottle is incubated and the colonies are counted and examined. This method gives better results than rinsing the bottle and subsequently plating the rinsings on the media. When used for this purpose, the agar concentration in Buffered Yeast Agar should be increased by 1% w/v (before sterilization).

Methodology

Suspend 41 grams of powder media in 1000 ml distilled water. Shake it well & heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 115°C for 20 minutes. Mix well and pour into sterile Petri plates.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Light amber coloured, clear to slightly opalescent gel forms in Petri plates.

Reaction

Reaction of 4.1% w/v aqueous solution at 25°C. pH : 5.5±0.2

pH range 5.30-5.70

Cultural Response/ characteristics

DM 1585: Cultural characteristics observed after an incubation at 25-30°C for 48-72 hours.

Organism	Inoculum (CFU)	Growth	Recovery
Candida albi cans ATCC 10231	50-100	good-luxuriant	>=70%
Saccharomyces cerevisiae ATCC 9763	50-100	good-luxuriant	>=70%
Aspergillus brasiliensis ATCC 16404	50-100	good-luxuriant	
Key *- Formerly known as Aspergillus niger			

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Ausubel, Brent, Kingston, Moore, Seidman, Smith and Struhl, 1994, Current Protocols in Molecular Biology, Current, Brooklyn, N.Y.
2. Davis J. G., 1931, J. Dairy Res., 3:133.
3. Bunker H. J., 1952, Lab. Prac., 18:354.
4. Bunker H. J., 1956, Wallerstein Lab. Communications, 19(65): 143.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Donot use the products if it fails to meet specifcatons for identity and performens parameters.