



Dehydrated Culture Media  
Bases / Media Supplements

## Technical Information

### Thioglycollate Agar

**Product Code: DM 1608**

**Application:** - Thioglycollate Agar is recommended for cultivation of anaerobic microorganisms.

### Composition\*\*

Ingredients	Gms / Litre
Casein enzymic hydrolysate	15.000
L-Cystine	0.500
Dextrose	5.500
Yeast extract	5.000
Sodium chloride	2.500
Sodium thioglycollate	0.500
Resazurin, sodium salt	0.001
Agar	20.000
Final pH ( at 25°C)	7.1±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Principle & Interpretation

In sterility tests thioglycollate Agar is used for the cultivation of aerobic as well as anaerobic microorganisms. It is based on the formula specified by US Pharmacopoeia <sup>(1)</sup> and APHA <sup>(2)</sup>. This medium neither requires anaerobic jar nor any special sealing for the cultivation of anaerobes. Thioglycollate Agar is also recommended for the cultivation of Clostridium species <sup>(1)</sup> and in the culture of Desulfotomaculum nigrificans.

Casein enzymic hydrolysate, yeast extract provides nitrogenous compounds, vitamin B and other essential growth nutrients. Dextrose is the fermentable carbohydrate and energy source. Resazurin is the redox indicator. Thioglycollate neutralizes the bacteriostatic effect of mercurial compounds used as the preservatives in the injection solution. If the solution used in test is a bacteriostatic ingredient then it is necessary to ascertain the bacteriostatic activity of the product.

### Methodology

Suspend 49.0 grams of powder media in 1000 ml distilled water. Shake well & heat to dissolve the medium completely. Dispense as desired and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Note: If more than the upper one-third has acquired a green colour, the medium may be restored once by heating in a waterbath or free flowing steam until the green colour disappears.

### Quality Control

#### Physical Appearance

Cream to yellow homogeneous free flowing powder

#### Gelling

Firm, comparable with 2.0% Agar gel.

#### Colour and Clarity of prepared medium

Light amber coloured clear to slightly opalescent (turning red due to aeration on standing) gel forms in Petri plates.

#### Reaction

Reaction of 4.9% w/v aqueous solution at 25°C. pH : 7.1±0.2

**pH range** 6.90-7.30



#### Cultural Response/ characteristics

DM1608: Cultural characteristics observed after an incubation at 35-37°C for 40-48 hours under anaerobic condition.

Organism	Inoculum (CFU)	Growth	Recovery
Clostridium botulinum ATCC 25763	50-100	luxuriant	>=50%
Clostridium perfringens ATCC 12924	50-100	luxuriant	>=50%
Clostridium sporogenes ATCC 11437	50-100	luxuriant	>=50%

#### Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

#### Further Reading

1. The United States Pharmacopoeia, 1985 21st rev. U.S. Pharmacopocial Convention, Rockville, M.D.
2. Speck M. L.(ed.), 1985, Compendium of Methods for the Microbiological examination of Foods, 2nd ed., APHA, Washington, D.C.

#### Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specificatons for identity and performens parameters.