

Technical Information

SPS Agar

Product Code: DM 1632

Application: Sulphite-Polymyxin-Sulphadiazine Agar is used for the detection of *Clostridium perfringens* in foods.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	15.000
Yeast extract	10.000
Sodium sulphite	0.500
Polymyxin B sulphate	0.010
Sulphadiazine	0.120
Ferric citrate	0.500
Agar	13.900
Final pH (at 25°C)	7.0±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Based on the Wilson and Blair Medium and the medium described by Mossel et al ^(2, 3) for selective isolation and enumeration of *C. perfringens* from foods. SPS (Sulphite Polymyxin Sulphadiazine) Agar was developed by Angelotti et al ⁽¹⁾. The medium of Mossel et al included the use of Miller-Prickett tubes. The modified SPS Agar however does not require the inclusion of Miller-Prickett tubes when used in the laboratory.

Casein enzymic hydrolysate and yeast extract supply nitrogenous compounds, vitamin B complex and other essential growth nutrients to the growing *C. perfringens*. This organism reduces sulphite to sulphide which reacts with iron of ferric citrate to form a black precipitate of iron sulphide and hence the colonies appear black ⁽⁴⁾. Polymyxin B and sulphadiazine inhibit a wide variety of gram-positive and gram-negative bacteria ⁽⁵⁾. Few organisms other than *C. perfringens* found in food also form black colonies on this medium.

Prepare serial dilutions of the samples to be tested and inoculate onto SPS Agar using the pour plate technique. If desired, pour cover layers using about 5 ml medium. Incubate the plates anaerobically. Enumerate the black colonies. Presumptive black *C. perfringens* colonies should be confirmed by additional standard biochemical tests.

Methodology

Suspend 40.03 grams of powder media in 1000 ml distilled water. Shake well & heat, to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.39% Agar gel

Colour and Clarity of prepared medium

Medium amber coloured slightly opalescent gel forms in Petri plates

Reaction

Reaction of 4.0% w/v aqueous solution at 25°C. pH : 7.0±0.2

pH range

6.80-7.20



Dehydrated Culture Media
Bases / Media Supplements

Cultural Response/Characteristics

DM 1632: Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours under anaerobic conditions.

Organism	Inoculum (CFU)	Growth	Recovery	Colour of colony
<i>Clostridium perfringens</i> ATCC 12924	50-100	Good-luxuriant	>=50%	Black
<i>Clostridium sporogenes</i> ATCC 11437	50-100	Fair-good	30-40%	Black
<i>Escherichia coli</i> ATCC 25922	50-100	inhibited	0%	
<i>Staphylococcus aureus</i> ATCC 25923	50-100	None-poor	<=10%	white

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Angelotti R., Han H. E., Foter M. J. and Lewis K. H., 1962, Appl. Microbiol., 10:193.
2. Mossel D. A. A., De Bruin A. S., Van Dipen H. M. J., Vendring C. M. A. and Zoutewelle G., 1956, J. Appl. Microbiol., 19:142.
3. Mossel R. S., 1959, J. Sci. Food Agric., 19:662.
4. Downes F. P. and Ito K., (Eds.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., APHA, Washington, D.C.
5. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams and Wilkins, Baltimore

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.

