

## Technical Information

### Milk Salt Agar Base

**Product Code: DM 1661**

**Application:** - Milk Salt Agar Base is used for selective isolation and cultivation of Staphylococci.

### Composition\*\*

Ingredients	Gms / Litre
Peptic digest of animal tissue	5.000
Beef extract	3.000
Sodium chloride	65.000
Agar	15.000
Final pH ( at 25°C)	7.4±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Principle & Interpretation

Milk Salt Agar is used for selective isolation and cultivation of Staphylococci. Koch reported that only Staphylococci could grow on agar media containing 7.5% sodium chloride <sup>(1)</sup>. Chapman in the modification of the Kochs medium utilized this property for making the medium more selective by using high salt content <sup>(2, 3)</sup>.

This is a simple but nutritious medium. Beef extract, peptic digest of animal tissue and skim milk supply essential nutrients mainly nitrogenous and carbonaceous compounds including trace ingredients to Staphylococci. Sodium chloride at a concentration of 6.5% makes the medium highly selective as majority of the contaminating organisms are inhibited by the high salt concentration, but Staphylococci are able to tolerate the high sodium chloride concentration.

### Methodology

Suspend 88 grams of powder media in 900 ml distilled water. Shake well & heat to dissolve the medium completely. Dispense and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 60°C. Aseptically add 10 ml of sterile skim milk (10% w/v skim milk powder solution) to every 90 ml of basal medium. Mix well and pour into sterile Petri plates.

### Quality Control

#### Physical Appearance

Off-white to yellow homogeneous free flowing powder

#### Gelling

Firm, comparable with 1.5% Agar gel

#### Colour and Clarity of prepared medium

Yellow coloured opaque gel forms in Petri plates after addition of 10%v/v sterile milk

#### Reaction

Reaction of the basal medium (8.8gm in 90 ml distilled water) at 25°C. pH : 7.4±0.2

#### pH range

7.20-7.60

#### Cultural Response/Characteristics

DM1661: Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours.



Dehydrated Culture Media  
Bases / Media Supplements

Organism	Inoculum (CFU)	Growth	Recovery
<i>Escherichia coli</i> ATCC 25922	>10 <sup>3</sup>	inhibited	0%
<i>Staphylococcus aureus</i> ATCC 25923	50-100	Good-Luxuriant	>=50%

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

**Prepared Media:** 2-8<sup>0</sup> in sealable plastic bags for 2-5 days.

## Further Reading

1. Koch, 1942, Zentralbl. Bakteriologie. Parasitenkunde. Abt. I. Orig., 149: 122.
2. Chapman, 1946, J. Bacteriology, 5 1:409.
3. Rechcigl M., Jr. (Ed.), 1978, CRC Handbook Series in Nutrition and Food, Section G., Vol. III, CRC Press, Inc., Ohio, U.S.A.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.

