

## Technical Information

### Moeller Decarboxylase Broth w/ Arginine HCl

#### Product Code: DM 1689

**Application:** - Moeller Decarboxylase Broth with Arginine hydrochloride, is used to differentiate bacteria on the basis of their ability to decarboxylate the L-Arginine hydrochloride.

#### Composition\*\*

Ingredients	Gms / Litre
Peptic digest of animal tissue	5.000
Beef extract	5.000
Dextrose	0.500
Bromocresol purple	0.010
Cresol red	0.005
Pyridoxal	0.005
L-Arginine hydrochloride	10.000
Final pH (25°C)	6.0±0.2

\*\*Formula adjusted, standardized to suit performance parameters

#### Principle & Interpretation

Moeller Decarboxylase Broth with Arginine hydrochloride is used for identification of gram-negative enteric bacilli on the basis of their ability to decarboxylate L-Arginine hydrochloride. Decarboxylase Broth was introduced by Moeller for detecting the production of lysine and ornithine decarboxylase and arginine dihydrolase <sup>(1)</sup>. Prior to Moellers work, bacterial amino acid decarboxylases were studied according to Gale <sup>(2)</sup> and Gale and Epps <sup>(3)</sup>. Decarboxylase media are also recommended by standard methods for identification of bacteria <sup>(4-7)</sup>.

This medium contains beef extract and peptic digest of animal tissue which provide nitrogenous nutrients for the growth of bacteria.

Dextrose is the fermentable carbohydrate and pyridoxal is the co-factor for the decarboxylase enzyme. Bromo cresol purple and cresol red are the pH indicators in this medium. When the medium is inoculated with dextrose fermenting bacteria, the pH is lowered due to acid production which changes the colour of the media from purple to yellow.

Acid produced stimulates decarboxylase enzyme. Arginine is first hydrolyzed to ornithine which is then decarboxylated to form putrescine. Formation of the amine putrescine increases the pH of the medium, changing the colour of the indicator from yellow to purple. If the organisms do not produce the appropriate enzyme, the medium remains acidic, yellow in colour. Each isolate to be tested should also be inoculated into the basal medium tube lacking the amino acid. After incubation, a decarboxylase test may show two layers of different colours, yellow and purple. Shake the tube gently before interpreting the results <sup>(8)</sup>.

Inoculated tubes must be protected from air with a layer of sterile mineral oil. Exposure to air may cause alkalization at the surface of the medium which makes the test invalid.

#### Methodology

Suspend 20.52 grams of powder media in 1000 ml distilled water. Shake well & heat if necessary to dissolve the medium completely.

Dispense in 5 ml amount in screw-capped tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 10 minutes.

## Quality Control

### Physical Appearance

Light yellow to greenish yellow homogeneous free flowing powder

### Colour and Clarity of prepared medium

Purple coloured clear solution without any precipitate

### Reaction

Reaction of 2.05% w/v aqueous solution at 25°C. pH : 6.0±0.2

**pH range** 5.80-6.20

### Cultural Response/ characteristics

DM 1689: Cultural characteristics observed after an incubation at 35-37°C for upto 4 days (Inoculated tubes are overlaid with sterile mineral oil).

Organism	Inoculum (CFU)	Ornithine decarboxylation
Citrobacter freundii ATCC 8090	50-100	variable reaction
Enterobacter aerogenes ATCC 13048	50-100	positive reaction, purple colour
Escherichia coli ATCC 25922	50-100	variable reaction
Klebsiella pneumoniae ATCC 13883	50-100	negative reaction, yellow colour
Proteus mirabilis ATCC 25933	50-100	negative reaction, yellow colour
Proteus vulgaris ATCC 13315	50-100	negative reaction, yellow colour
Pseudomonas aeruginosa ATCC 9027	50-100	positive reaction, purple colour
Salmonella Paratyphi A ATCC 9150	50-100	delayed positive reaction/ positive reaction, purple colour
Salmonella Typhi ATCC 6539	50-100	delayed positive reaction/ negative reaction, yellow colour
Serratia marcescens 8100	50-100	negative reaction, yellow colour
Shigella dysenteriae ATCC 13313	50-100	delayed positive reaction/ negative reaction, yellow colour
Shigella flexneri ATCC 12022	50-100	delayed positive reaction/ negative reaction, yellow colour
Shigella sonnei ATCC 25931	50-100	variable reaction

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

## Further Reading

1. Moeller V., 1955, Acta Pathol. Microbiol. Scand. 36:158.
2. Gale G. F., 1940, Biochem. J., 34:392.
3. Gale and Epps, 1943, Nature, 152:327.
4. Isenberg (Ed.), 1992, Clinical Microbiology Procedures Handbook, Vol. I, ASM, Washington, D. C.
5. FDA Bacteriological Analytical Manual, 8th Ed., AOAC International, Gaithersburg, Md.
6. Eaton A. D., Clesceri L. S. and Greenberg A. E., (Ed.), 1998, Standard Methods for the Examination of Water and Wastewater, 20th Ed., American Public Health Association, Washington, D.C
7. Downes F. P. and Ito K., (Ed.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., American Public Health Association, Washington, D.C
8. MacFaddin J. F., 2000, Biochemical tests for Identification of Medical Bacteria, 3rd Ed., Lippincott, Williams and Wilkins, Baltimore.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Donot use the products if it fails to meet specificatons for identity and performens parameters.