

Technical Information

Ascospore Agar

Product Code: DM 1804

Application: - Ascospore Agar is recommended for enrichment and detection of ascosporegenous yeasts.

Composition**

Ingredients	Gms / Litre
Yeast extract	2.500
Dextrose	1.000
Potassium acetate	10.000
Agar	30.000
Final pH (at 25°C)	6.4±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Ascospore Agar is recommended for the enrichment and detection of ascospores in ascosporegenous yeasts such as *Saccharomyces cerevisiae*. It is based on the formula developed by McClary et al ⁽¹⁾. Ascospore Agar is the modification of McClary medium with the addition of potassium acetate in place of sodium acetate. Acetate salt of potassium was found to be superior to sodium salt for sporulation in *Saccharomyces* ^(1, 2).

Dextrose and yeast extract provide the nutrients required for the growth and also stimulate ascospore formation in yeasts. Slightly acidic pH of the medium favours the growth of *Saccharomyces cerevisiae*.

Methodology

Suspend 43.50 grams of powder media in 1000 ml distilled water. Shake well & heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.

Quality Control

Physical Appearance

Light yellow to brownish yellow homogeneous free flowing powder

Gelling

Firm, comparable with 3.0% Agar gel

Colour and Clarity of prepared medium

Medium amber coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 4.35% w/v aqueous solution at 25°C. pH : 6.4±0.2

pH range 6.20-6.60

Cultural Response/Characteristics

DM1804: Cultural characteristics observed after an incubation at 25-30°C for upto 3-6 days.

Organism	Inoculum (CFU)	Growth	Recovery	Ascospores
<i>Candida albicans</i> ATCC 10231	50-100	luxuriant	>=50%	negative
<i>Saccharomyces cerevisiae</i> ATCC 9763	50-100	luxuriant	>=50%	positive



Dehydrated Culture Media
Bases / Media Supplements

Storage and Shelf Life

Dried media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8⁰ in sealable plastic bags for 2-5 days.

Further Reading

1. McClary D.O., Nulty W.L. and Miller G.R., 1959, J.Bacteriol., 78:362

2. MacFaddin J.F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Williams and Wilkins, Baltimore.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.

