

Technical Information

Vibrio Agar

Product Code: DM 1820

Application: - Vibrio Agar is used for selective cultivation of *Vibrio* species.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	4.000
Yeast extract	5.000
Proteose peptone	3.000
Sucrose	20.000
Sodium thiosulphate	6.500
Sodium citrate	10.000
Sodium deoxycholate	1.000
Sodium chloride	10.000
Oxgall	5.000
Sodium lauryl sulphate	0.200
China blue	0.200
Cresol red	0.020
Agar	15.000
Final pH (at 25°C)	8.5±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Vibrio species, like many other gram-negative bacteria, can grow in the presence of high levels of bile salts. They are facultatively anaerobic and grow best in alkaline conditions. Isolation is facilitated by the use of media formulated with an alkaline pH and selective for Vibrios by adding appropriate selective agents. The main agents employed are bile salts, teepol, tellurite and polymyxin B and E (Colistin) ⁽²⁾.

Vibrio Agar is a selective medium for the isolation of *Vibrio cholerae*, *Vibrio parahaemolyticus* and other *Vibrios* ⁽¹⁾.

Casein enzymic hydrolysate, proteose peptone, yeast extract provide nitrogenous, carbonaceous compounds, sulphur, vitamin B complex and other essential growth nutrients. Sodium citrate, sodium deoxycholate and oxgall inhibit gram-positive organisms and coliforms. Sucrose is the fermentable carbohydrate which is fermentative by *V. cholerae* and *V. alginolyticus* and form blue colonies due to the indicator china blue. *V. parahaemolyticus* forms slightly reddish and translucent colonies. Sodium thiosulphate in combination with ferric citrate detects H₂S production. Thiosulphate also acts as a sulphur source. Alkaline pH of this medium helps in recovery of *V. cholerae*. China blue and cresol red are the pH indicators.

Methodology

Suspend 79.92 grams of powder media in 1000 ml distilled water. Shake well & heat to dissolve the medium completely. DO NOT AUTOCLAVE. Cool to 45-50°C and pour into sterile Petri plates.

Quality Control

Physical Appearance

Light yellow to greyish yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel.

Colour and Clarity of prepared medium

Reddish purple coloured clear to slightly opalescent gel forms in Petri plates.

Reaction

Reaction of 7.99% w/v aqueous solution at 25°C. pH : 8.5±0.2

pH Range:-

8.30-8.70

Cultural Response/Characteristics

DM 1820: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum(CFU)	Growth	Recovery	
<i>Enterococcus faecalis</i> ATCC 29212	50-100	none-poor	<=10%	yellow
<i>Escherichia coli</i> ATCC 25922	>=10 ³	inhibited	0%	-
<i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	none-poor	<=10%	blue
<i>Salmonella Typhi</i> ATCC 6539	>=10 ³	inhibited	0%	-
<i>Shigella flexneri</i> ATCC 12022	>=10 ³	inhibited	0%	-
<i>Vibrio cholerae</i> ATCC 15748	50-100	good-luxuriant	>=50%	blue
<i>Vibrio parahaemolyticus</i> ATCC 17802	50-100	good-luxuriant	>=50%	slightly reddish

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8⁰ in sealable plastic bags for 2-5 days.

Further Reading

1. Atlas R. M. 2004, 3rd Ed., Handbook of Microbiological Media, Parks, L.C., (Ed.), CRC Press, Boca Raton.
2. Gomez-Gil B. and Roque A., Isolation, Enumeration and Preservation of the Vibrionaceae, Thompson F. L., Austin B. and Swings J., The Biology of Vibrios, ASM press.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Donot use the products if it fails to meet specificatons for identity and performens parameters.