

Technical Information

M2 Agar

Product Code: DM 1858

Application: - M2 Agar is a semi synthetic culture medium used as a general purpose plate count agar.

Composition**

Composition		
Ingredients	Gms / Litre	
Yeast extract	5.000	
Dextrose	10.000	
Sodium chloride	10.000	
Monopotassium phosphate	0.100	
Magnesium sulphate	0.050	
Agar	15.000	
Final pH (at 25°C)	7.1±0.2	
**Formula adjusted, standardized to suit performance par	ameters	

Principle & Interpretation

Nutrient media are basic culture media used for maintaining & cultivating microorganisms, including fastidious organisms ^{(1, 2).} Plate Count Agar is recommended for the count of microorganisms in food, dairy products, water and waste water. M2 Agar is suitable for determining bacterial count in sterile rooms.

Yeast extract provides a source of trace elements, vitamins and amino acids. Dextrose is the source of carbohydrate. Monopotassium phosphate buffers the media. Magnesium sulphate is a source of divalent cations. Sodium chloride is an essential ion and helps in maintaining the osmotic balance of the medium. Agar is the solidifying agent.

Methodology

Suspend 40.15 grams of powder media in 1000 ml distilled water containing 132 ml glycerine. Shake well & heat to dissolve the medium completely. sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Light amber coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 4.015% w/v aqueous solution at 25°C. pH: 7.1±0.2

pH range 6.90-7.30

Cultural Response/Characteristics

DM1858: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

OrganismGrowthSaccharomyces cerevisiae ATCC 9763Good-luxuriantEscherichia coli ATCC25922Good-luxuriant





Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label. **Prepared Media:** 2-8⁰ in sealable plastic bags for 2-5 days.

Further Re	eading
------------	--------

	bbons D., (Eds.), Vol. 3A, Academic Press, London.	, Norris J. and Ribbons D., (s in Microbiology'	J. and Mitchell T., 1970, Metho	1. Lapage S., Shelton J.
--	--	-------------------------------	--------------------	---------------------------------	--------------------------

2. MacFaddin J. F 2000. Biochemical 🛚	facts for Idontification of Modical D	actoria 2rd Ed Linnincott	Williams and Wilkins Paltimore
2. Macrauulli J. F., 2000. Biochelliicai	esis idi idelililicalidii di Medicai b	acteria. Siu Lu., Libbilicott.	Williams and Wilkins, Baltimore.

Disclaimer:

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performens parameters.

