

## Technical Information

### Decarboxylase Test Medium Base (Falkow)

#### Product Code: DM 1912S

**Application:** - Decarboxylase Test Medium Base (Falkow) is used for testing decarboxylase activity.

#### Composition\*\*

Ingredients	Gms / Litre
Peptic digest of animal tissue	5.000
Yeast extract	3.000
Dextrose	1.000
Bromo cresol purple	0.020
Final pH ( at 25°C)	6.7±0.2

\*\*Formula adjusted, standardized to suit performance parameters

#### Principle & Interpretation

First practical application of amino acid decarboxylase test was reported by Moeller for distinguishing various microorganisms<sup>(1)</sup>. Moellers work was based on the earlier experiments done by Gale<sup>(2)</sup> and Gale and Epps<sup>(3)</sup> on bacterial amino acid decarboxylases. He observed that production of lysine, arginine, ornithine decarboxylase by various members of *Enterobacteriaceae* provide an important parameter to including other biochemical tests for differentiating bacteria within closely related groups. Further, to differentiate *Salmonella arizonae* from *Citrobacter*, Calquist<sup>(4)</sup> developed a medium utilizing the lysine decarboxylase reaction. Later on Falkow<sup>(5)</sup> developed the lysine decarboxylase medium for differentiating *Salmonellae* and *Shigellae* by the valid and reliable results. This medium is also recommended by BIS for detection of dihydrolase and decarboxylase activity of *Vibrio cholerae* and other vibrios<sup>(6)</sup>.

Dextrose is fermented by the enteric bacteria resulting in acidic pH. Bacteria which produce lysine or ornithine or arginine decarboxylase after 24-96 hours will indicate an alkaline reaction seen as purple colour for decarboxylase producing bacteria and an acid pH (yellow) by the bacteria not producing decarboxylase. Inoculated tubes must be protected from air (by overlaying the medium with sterile mineral oil) to avoid false alkalization at the surface of the medium. Control tubes of basal media should be inoculated.

The decarboxylase reactions can be considered indicative of a given genus or species but conclusive and final identification of these organisms cannot be made solely on the basis of the decarboxylase reactions.

#### Methodology

Suspend 9 grams of powder media in 1000 ml distilled water. Shake well & heat to dissolve the medium completely. Divide into four equal parts. One part is tubed without addition of any amino acid. To the remaining three parts, add separately L-lysine hydrochloride, L-arginine hydrochloride and L-ornithine hydrochloride to a final concentration of 0.5%. Dispense in 3-4 ml quantities in screw capped tubes and sterilize by autoclaving at 10 lbs pressure (115°C) for 20 minutes. To avoid false alkalization at the surface of medium it is recommended to add liquid paraffin to a height of about 5mm before sterilization.

#### Quality Control

##### Physical Appearance

Yellow to greenish yellow coloured homogeneous free flowing powder

##### Colour and Clarity of prepared medium

Purple coloured clear solution without any precipitate.

##### Reaction

Reaction of 0.9% w/v aqueous solution at 25°C. 6.7±0.2

pH range : 6.5-6.9

##### Cultural Response/ characteristics

DM 1912S: Cultural characteristics observed after an incubation at 35-37°C for upto 4 days.



Dehydrated Culture Media  
Bases / Media Supplements

Organism	Lysine Decarboxylation	Arginine Decarboxylation	Ornithine Decarboxylation
<i>Enterobacter aerogenes</i> ATCC 13048	positive reaction, purple colour	negative reaction, yellow colour	positive reaction, purple colour
<i>Escherichia coli</i> ATCC 25922	variable reaction	variable reaction	variable reaction
<i>Klebsiella pneumoniae</i> ATCC 13883	positive reaction, purple colour	negative reaction, yellow colour	negative reaction, yellow colour
<i>Proteus vulgaris</i> ATCC 13315	negative reaction, yellow colour	negative reaction, yellow colour	negative reaction, yellow colour
<i>Pseudomonas aeruginosa</i> ATCC 27853	negative reaction, yellow colour	positive reaction, purple colour	negative reaction, yellow colour
<i>S. serotype typhi</i> ATCC 6539	positive reaction, purple colour	delayed positive reaction or negative reaction, yellow colour	negative reaction, yellow colour
<i>Serratia marcescens</i> ATCC 8100	positive reaction, purple colour	negative reaction, yellow colour	positive reaction, purple colour
<i>Shigella flexneri</i> ATCC 12022	negative reaction, yellow colour	delayed positive reaction or negative reaction, yellow colour	negative reaction, yellow colour
<i>Vibrio cholerae</i> ATCC 15748	negative reaction, yellow colour	positive reaction, purple colour	positive reaction, purple colour

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

## Further Reading

1. Moeller, 1954, Acta Path. Micro. Scand., 34:102.
2. Gale, 1940, Biochem. J., 34:392, 583, 846.
3. Gale and Epps, 1943, Nature, 152:327.
4. Calquist, 1956, J. Bact., 71:339.
5. Falkow, 1958, Am. J. Clin. Path., 29:598.
6. Bureau of Indian Standards, IS : 5887 (Part V) 1976, reaffirmed 1986.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

