

Technical Information

PSTA Enrichment Broth Base

Product Code: DM 1940

Application: - PSTA Enrichment Broth Base is recommended for secondary enrichment of *Yersinia enterocolitica* from foods.

Composition**

Ingredients	Gms / Litre
Peptic digest of animal tissue	1.000
Sucrose	1.000
Tris hydroxymethyl aminomethane	3.000
Brilliant green	0.0125
Sodium azide	0.192
Final pH (at 25°C)	8.3±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Yersinia enterocolitica are ubiquitous in nature being isolated frequently from soil, water, animals, and a different types of foods. They comprise a biochemically heterogeneous group that can grow at refrigeration temperatures. *Y. enterocolitica* has been isolated from environmental and food sources, such as ponds, lakes, meats, ice cream, and milk. *Yersinia* species have the ability to grow at 4°C⁽¹⁾ and also exhibit tolerance to dilute alkali^(2, 3). PSTA Enrichment Broth Base formulated in accordance with APHA⁽⁴⁾, is also recommended for secondary or selective enrichment of *Y. enterocolitica* which is advantageous as it provides higher selectivity, thereby increasing the chance of recovery or isolation of target organism. Peptic digest of animal tissue in the medium provides nitrogen, vitamins and minerals necessary to support bacterial growth. Sucrose is the carbohydrate source. Brilliant green and sodium azide inhibits the growth of gram-negative organisms. About 25 grams of food sample is added to 225 ml of PSB Broth Base (DM1941) and incubated at 4°C for 14-28 days. 1 ml of this primary enrichment is inoculated in 100 ml of PSTA Enrichment Broth Base (DM1940) and incubated at 28°C for 48 hours. The secondary enrichment is then streaked on selective media such as SS Agar (DM1108), *Yersinia* Selective Agar Base (DM1843).

Methodology

Suspend 5.2 grams of powder media in 1000 ml distilled water. Shake well & sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C and aseptically add Ampicillin to a final concentration 0.005 gm per litre. Mix well and dispense as desired.

Quality Control

Physical Appearance

Light yellow to greenish yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Green coloured clear to slightly opalescent solution

Reaction

Reaction of 0.52% w/v aqueous solution at 25°C. pH : 8.3±0.2

pH Range 8.10-8.50

Cultural Response/ characteristics

DM 1940: Cultural characteristics observed after an incubation at 28°C for 48 hours.

Organism	Inoculum (CFU)	Growth
<i>Yersinia enterocolitica</i> ATCC 27729	50-100	good-luxuriant



Dehydrated Culture Media
Bases / Media Supplements

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Pai C. H. and Mors V., 1978, Infect. Immun., 19: 908-911
2. Aulisio C. C. G., Mehlman I. J. and Sander A. C., 1980, Appl. Environ. Microbiol. 39: 135-140
3. Doyle M. P. and Hugdahl M. B., 1983, Appl. Environ. Microbiol., 45:127-135
4. Speck M. L., (Eds.), 1984, Compendium of Methods for the Microbiological Examination of Foods, 2nd Ed., APHA, Washington, D.C.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Donot use the products if it fails to meet specificatons for identity and performens parameters.

