



Dehydrated Culture Media
Bases / Media Supplements

Technical Information

Tryptose Broth, w/ Thiamine HCl

Product Code: DM 1997

Application: - Tryptose Broth, w/Thiamine HCl is used for the cultivation and differentiation of fastidious microorganisms in an infusion free medium.

Composition**

Ingredients	Gms / Litre
Tryptose	20.000
Dextrose	1.000
Sodium chloride	5.000
Thiamine hydrochloride	0.005
Final pH (at 25°C)	7.2±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Tryptose containing media, rather than the conventionally used meat infusion media have been used for the enumeration and isolation of *Brucella* species^(2,3). Addition of thiamine to tryptose media enhanced the recovery of *Brucella* species especially *Brucella suis*^(4,5).

This medium can be used as general purpose medium for cultivation of wide variety of organisms. It can also be supplemented with defibrinated blood (sheep, horse) to prepare blood containing medium for the isolation of fastidious organisms like *Brucella*. Tryptose Broth with thiamine HCl is recommended by APHA⁽⁶⁾ and Diagnostic Procedures and Reagents⁽⁷⁾ for the isolation and cultivation of *Brucella* species including Streptococci, meningococci, pneumococci and other pathogenic bacteria⁽⁸⁾.

Dextrose is the source of energy. Tryptose serves as nitrogen source while sodium chloride maintains osmotic equilibrium.

Methodology

Suspend 26 grams of powder media in 1000 ml distilled water. If desired, add 0.5-1% agar to the medium. Shake well & heat to dissolve the media completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. For blood media, aseptically add 5% v/v sterile defibrinated blood. Mix well and dispense as desired.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Basal Medium: Yellow coloured clear to slightly opalescent Solution. After addition of 5% v/v sterile defibrinated blood: cherry red coloured opaque solution forms in tubes.

Reaction

Reaction of 2.6% w/v aqueous solution at 25°C. pH : 7.2±0.2

pH Range:-

7.00-7.40

Cultural Response/Characteristics

DM 1997: Cultural characteristics observed after an incubation at 35-37°C for 48-72 hours with added 5% v/v sterile defibrinated blood in presence of 10% Carbon dioxide (CO₂).





Dehydrated Culture Media
Bases / Media Supplements

Organism	Inoculum (CFU)	Growth
<i>Brucella melitensis</i> ATCC 4309	50-100	good-luxuriant
<i>Brucella suis</i> ATCC 4314	50-100	good-luxuriant
<i>Streptococcus pneumoniae</i> ATCC 6303	50-100	good-luxuriant
<i>Streptococcus pyogenes</i> ATCC 19615	50-100	good-luxuriant

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8^o in sealable plastic bags for 2-5 days.

Further Reading

1. Huddleson I. F., 1943, Brucellosis in man and animals, rev., Ed., The Commonwealth Fund, New York, N.Y.
2. Ruiz Castañeda M., 1947, Proc. Soc. Exp. Biol. Med., 64:114.
3. Huddleson I. F., 1939, Brucellosis in Man and Animals, Oxford University Press, Oxford, England.
4. McCullough W. G., Mills R. L., Herbst E. J., Roessler W. J. and Brewer C. R., 1947, J. Bacteriol., 53:5.
5. Atlas R. M., 2004, Handbook of Microbiological Media, Lawrence C. Parks, (Ed.), 3rd Edition, CRC Press.
6. Standard Methods for the Microbiological Examination of Dairy Products, 9th Ed., 1948, APHA Inc., New York.
7. Diagnostic Procedures and Reagents, 1950, 3rd Edition, APHA, New York.
8. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams and Wilkins, Baltimore

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.

