

Dehydrated Culture Media Bases / Media Supplements

# **Technical Information**

## **Phenol Red Xylose Broth**

## Product Code: DM2015

Application: - Phenol Red Xylose Broth is used for xylose fermentation studies of microorganisms.

Composition**				
Ingredients	Gms / Litre			
Proteose peptone	10.000			
Beef extract	1.000			
Sodium chloride	5.000			
Xylose	5.000			
Phenol red	0.018			
Final pH ( at 25°C)	7.4±0.2			
**Formula adjusted, standardized to suit per	ormance parameters			

### **Principle & Interpretation**

Phenol Red Broth Medium is formulated by Vera<sup>(2)</sup> is recommended to determine the fermentation reaction of different carbohydrates for the identification of microorganisms<sup>(3-5)</sup>. Phenol Red Broth Medium with various carbohydrates serves as a differential medium by helping in differentiation of various species and genera by their ability to ferment the specific carbohydrate, with the production of acid or acid and gas<sup>(6)</sup>. Phenol Red Xylose Broth is used to study xylose fermentation in various bacteria.

Proteose peptone and beef extract serve as sources for carbon and nitrogen. Sodium chloride is the osmotic stabilizer. Phenol red is the pH indicator, which turns yellow at acidic pH i.e. on fermentation of xylose. Gas formation is seen in Durhams tubes. All the members of family *Enterobacteriaceae* grow well in this medium. In addition to producing a pH colour shift, the production of mixed acids, notably butyric acids, often results in a pungent, foul odour from the culture medium <sup>(1)</sup>.

## Methodology

Suspend 21 grams of powder media in 1000 ml distilled water and mix well. Shake well & heat if necessary to ensure complete

dissolution. Distribute in fermentation tubes (tubes containing inverted Durham's tubes). Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

## Quality Control

Physical Appearance

Light yellow to pink coloured homogeneous free flowing powder

#### Colour and Clarity of prepared medium

Red coloured Clear solution without any precipitate

Reaction

Reaction of 2.1% w/v aqueous solution at 25°C. pH : 7.4±0.2

pH Range 7.20-7.60

#### Cultural Response/ characteristices

DM 2015: Cultural characteristics observed after an incubation at 35 - 37°C for 18 - 24 hours.





Dehydrated Culture Media Bases / Media Supplements

Organism	Inoculum (CFU)	Growth	Acid
Citrobacter freundii ATCC 8090	50-100	luxuriant	Positive reaction, yellow colour
Escherichia coli ATCC 25922	50-100	luxuriant	Positive reaction, yellow colour
Enterobacter aerogenes ATCC 13048	50-100	luxuriant	Positive reaction, yellow colour
Klebsiella pneumoniae ATCC 13883	50-100	luxuriant	Positive reaction, yellow colour
Proteus vulgaris ATCC 13315	50-100	luxuriant	Positive reaction, yellow colour
Salmonella Typhi ATCC 6539	50-100	luxuriant	Positive reaction, yellow colour
Salmonella Typhimurium ATCC 1402	850-100	luxuriant	Positive reaction, yellow colour
Serratia marcescens ATCC 8100	50-100	luxuriant	Negative reaction, no colour change
Shigella flexneri ATCC 12022	50-100	luxuriant	Negative reaction, no colour change

## Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label. Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

### **Further Reading**

1. Koneman E. W., Allen S. D., Janda W.M., Schreckenberger P.C., Winn W.C. Jr., 1992, Colour Atlas and Textbook of Diagnostic

Microbiology, 4th Ed., J. B. Lippinccott Company

2. Vera H. D., 1950, Am. J. Public Health, 40, 1267

3. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification -Maintenanceof Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.

4. Finegold S. M. and Baron E. J., 1986, Bailey and Scotts Diagnostic Microbiology, 7th Ed., The C.V. Mosby Co., St. Louis.

5. Ewing W. H., 1986, Edwards and Ewings Identification of Enterobacteriaceae, 4th ed.,Elsevier Science Publishing Co., Inc., New York. 6. MacFaddin J. F., 2000, Biochemical tests for Identification of Medical Bacteria, 3rd edi., Lippincott, Williams and Wilkins, Baltimore.

### **Disclaimer**:

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Donot use the products if it fails to meet specificatons for identity and performens parameters.



#### Gas

Positive reaction Positive reaction Positive reaction Weak reaction Negative reaction Negative reaction Negative reaction Negative reaction