

Technical Information

Phenol Red Inositol Broth

Product Code: DM 2017

Application: - Phenol Red Inositol Broth is used for inositol fermentation studies of microorganisms.

Composition**

Ingredients	Gms / Litre
Proteose peptone	10.000
Beef extract	1.000
Sodium chloride	5.000
Inositol	5.000
Phenol red	0.018
Final pH (at 25°C)	7.4±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Phenol Red Broth Medium is formulated by Vera ⁽²⁾ is recommended to determine the fermentation reaction of different carbohydrates for the identification of microorganisms ⁽³⁻⁵⁾. Phenol Red Broth Medium with various carbohydrates serves as a differential medium by helping in differentiation of various species and genera by their ability to ferment the specific carbohydrate, with the production of acid or acid and gas ⁽⁶⁾. Phenol Red Inositol Broth is used to study inositol fermentation in various bacteria.

Proteose peptone and beef extract serve as sources for carbon and nitrogen. Sodium chloride is the osmotic stabilizer. Phenol red is the pH indicator, which turns yellow at acidic pH i.e. on fermentation of inositol. Gas formation is seen in Durhams tubes. All the members of *Enterobacteriaceae* family grow well in this medium. In addition to producing a pH colour shift, the production of mixed acids, notably butyric acids, often results in a pungent, foul odour from the culture medium ⁽¹⁾.

Methodology

Suspend 21 grams of powder media in 1000 ml distilled water and mix well. Shake well and heat if necessary to ensure complete dissolution. Distribute in fermentation tubes (tubes containing inverted Durham's tubes). Sterilize by autoclaving at 15 lbs pressure (12 1°C) for 15 minutes.

Quality Control

Physical Appearance

Light yellow to pink coloured homogeneous free flowing powder

Colour and Clarity of prepared medium

Red coloured Clear solution without any precipitate

Reaction

Reaction of 2.1% w/v aqueous solution at 25°C. pH : 7.4±0.2

pH Range 7.20-7.60

Cultural Response/ characteristics

DM 2017: Cultural characteristics observed after an incubation at 35 - 37°C for 18 - 24 hours.



Dehydrated Culture Media
Bases / Media Supplements

Organism	Inoculum (CFU)	Growth	Acid	Gas
Citrobacter freundii ATCC 8090	50-100	luxuriant	Negative reaction, no colour change	Negative reaction
Escherichia coli ATCC 25922	50-100	luxuriant	Negative reaction, no colour change	Negative reaction
Enterobacter aerogenes ATCC 13048	50-100	luxuriant	Positive reaction, yellow colour	Positive reaction
Klebsiella pneumoniae ATCC 13883	50-100	luxuriant	Positive reaction, yellow colour	Positive reaction
Proteus vulgaris ATCC 13315	50-100	luxuriant	Negative reaction, no colour change	Negative reaction
Salmonella Typhi ATCC 6539	50-100	luxuriant	Negative reaction, no colour change	Negative reaction
Salmonella Typhimurium ATCC 14028	50-100	luxuriant	Positive reaction, yellow colour	Positive reaction
Serratia marcescens ATCC 8100	50-100	luxuriant	Positive reaction, yellow colour	Positive reaction
Shigella flexneri ATCC 12022	50-100	luxuriant	Negative reaction, no colour change	Negative reaction

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Koneman E. W., Allen S. D., Janda W.M., Schreckenberger P.C., Winn W.C. Jr., 1992, Colour Atlas and Textbook of Diagnostic Microbiology, 4th Ed., J. B. Lippincott Company
2. Vera H. D., 1950, Am. J. Public Health, 40, 1267
3. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification -Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore.
4. Finegold S. M. and Baron E. J., 1986, Bailey and Scotts Diagnostic Microbiology, 7th Ed., The C.V. Mosby Co., St. Louis.
5. Ewing W. H., 1986, Edwards and Ewings Identification of Enterobacteriaceae, 4th ed., Elsevier Science Publishing Co., Inc., New York.
6. MacFaddin J. F., 2000, Biochemical tests for Identification of Medical Bacteria, 3rd ed., Lippincott, Williams and Wilkins, Baltimore.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Donot use the products if it fails to meet specificatons for identity and performens parameters.

