

Technical Information

Antibiotic Assay Medium No. 41

Product Code: DM 2144

Application: - Antibiotic Assay Medium No. 41 is used for the microbiological assay of Thiostrepton using *Enterococcus hirae* as the test organism.

Composition**			
Ingredients	Gms / Litre		
Casein enzymic hydrolysate	9.000		
Dextrose	20.000		
Yeast extract	5.000		
Sodium citrate	10.000		
Monopotassium phosphate	1.000		
Dipotassium hydrogen phosphate	1.000		
Final pH (at 25°C)	6.8±0.2		
**Formula adjusted, standardized to suit performance	e parameters		

Principle & Interpretation

Antibiotic Assay Medium No. 41 is used for turbidimetric microbiological assay of thiostreptone, a polypeptide antibiotic.Grove and Randall have elucidated the antibiotic assays and media in their comprehensive treatise on antibiotic assays. (1)

Essential amino acids, mineral and growth factors are supplied by casein enzymic hydrolysate and yeast extract in this medium. Dextrose provides carbon and energy source for enhancing the growth of test organism. Good buffering action is maintained by phosphates in the medium. Sodium citrate provides additional source of carbon and energy and promote enhanced growth of the test organism.

Turbidimetric antibiotic assay is based on the change or inhibition of growth of a test microorganims in a liquid medium containing a uniform concentration of an antibiotic. After incubation of the test orgainism in the working dilutions of the antibiotics, the amount of growth is determined by measuring the light transmittance using spectophotometer. The concentration of antibiotic is determined by comparing amounts of growth obtained with that is given by the reference standard solutions. Use of this method is appropriate only when test samples are clear.

Methodology

Suspend 46.0 grams of powder media in 1000 ml distilled water. Shake well and heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.





Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Light yellow coloured clear solution

Reaction

Reaction of 4.6% w/v aqueous solution at 25°C. pH: 6.8±0.2

pH Range:- 6.60-7.00

Growth Promotion Test

In accordance with the harmonized method of USP

Cultural Response/Characteristics

DM2144: Cultural characteristics observed after an incubation at 35 - 37°C for 18 - 24 hours.

Organism	Inoculum (CFU)	Growth	Serial dilution with
Enterococcus hirae ATCC 10541	50-100	luxuriant	Thiostrepton

Storage and Shelf Life

Dried media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Grove and Randall, 1955; Assay methods of Antibiotics Medical Encyclopedia, Inc, New York.

Disclaimer:

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Donot use the products if it fails to meet specifications for identity and performens parameters.

