

Technical Information

Modified Duncan Strong (DS) Medium

Product Code: DM 2237

Application: - Modified Duncan Strong (DS) Medium is used for isolation and differentiation of *Clostridium perfringens* from other clostridia from foods on the basis of raffinose fermentation.

Composition**

Ingredients	Gms / Litre
Proteose peptone	15.000
Yeast extract	4.000
Sodium thioglycollate	1.000
Disodium phosphate	10.000
Raffinose	4.000
Final pH (25°C)	7.8±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Clostridium perfringens, a gram-positive, rod shaped, anaerobic, spore-forming bacteria, is the major cause of food poisoning in humans. This organism is commonly found in raw meats, poultry, dehydrated soups and sauces, raw vegetables and certain other foods or food ingredients.

A heat labile enterotoxin produced only by sporulating cells induces the major symptoms of diarrhea in perfringens poisoning.

Modified Duncan Strong (DS) Medium is formulated as per Duncan and Strong ⁽¹⁾, This is also recommended by APHA ⁽²⁾ for the isolation and differentiation of *C. perfringens* from other Clostridia from foods on the basis of raffinose fermentation. It is also used for the rapid detection of the *Clostridium perfringens* enterotoxin ⁽³⁾.

Proteose peptone and yeast extract provide nitrogenous compounds and other nutrients for bacterial growth. Sodium thioglycollate helps to create anaerobic conditions suitable for clostridial growth. Disodium phosphate acts as a buffering agent. Raffinose in the medium is fermented only by *C. perfringens* to produce acid within 72 hours of inoculation which can be tested by transfer 1 ml of culture to a test tube or spot plates and adding 2 drops of 0.04% bromothymol blue. A yellow colour indicates acid production.

Inoculate about 2 gm of the food sample into 15- 20 ml of Chopped Liver Broth (DM1606). After an incubation at 35-37°C for 20-24 hours, tubes showing turbidity are streaked on Perfringens Agar Base (DM1837) containing Egg Yolk Emulsion (MS2045) to obtain presumptive *C. perfringens*. These presumptive colonies can be confirmed by inoculating into Motility Nitrate Medium, Buffered (DM1630), Lactose Gelatin Medium (DM1628) and Modified Duncan Strong (DS) Medium (DM2237) (2).

Methodology

Suspend 34 grams of powder media in 1000 ml distilled water and mix thoroughly. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Dispense into sterile tubes. Check one or two tubes for measuring the pH.



Dehydrated Culture Media
Bases / Media Supplements

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Yellow coloured clear solution without any precipitate

Reaction

Reaction of 3.4% w/v aqueous solution at 25°C. pH : 7.8±0.2

pH range 7.60-8.00

Cultural Response/ characteristics

DM 2237: Cultural characteristics observed after an incubation at 35-37°C for 48-72 hours.

Organism	Inoculum (CFU)	Growth	Raffinose fermentation
Clostridium perfringens ATCC 12924	50-100	good-luxuriant	positive reaction
Clostridium sporogenes ATCC 11437	50-100	good-luxuriant	negative reaction

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Duncan C. and Strong D., 1969, Appl. Microbiol., 16: 82.
2. Labbe R. G. and Rey D. K., 1979, Appl. Microbiol., 13: 559.
3. Downes F. P. and Ito K., (Eds.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., APHA, Washington, D.C.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Donot use the products if it fails to meet specificatons for identity and performens parameters.

