

Technical Information

Sabouraud Dextrose Maltose Agar

Product Code: DM 2313

Application: - Sabouraud Dextrose Maltose Agar is used for the cultivation of molds, yeasts and aciduric organisms as well as testing antimycotic substances.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	5.000
Peptic digest of animal tissue	5.000
Dextrose	10.000
Maltose	10.000
Agar	15.000
Final pH (at 25°C)	5.4±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Sabouraud Dextrose Agar is Carliers modifications ⁽¹⁾ of the formulation described by Sabouraud ⁽²⁾ for the cultivation of fungi, particularly those associated with skin infections. Sabouraud Dextrose Maltose Agar is used for the cultivation of yeast, moulds and other aciduric organisms ^(3, 4, 5).

Sabouraud dextrose media are peptone media supplemented with dextrose to support the growth of fungi. Casein enzymic hydrolysate and peptic digest of animal tissue provide nitrogen, vitamins, minerals, amino acids and growth factors. Dextrose and maltose act as an energy source for the growth of microorganisms. The low pH favours fungal growth and inhibits contaminating bacteria from clinical specimens to grow ⁽⁶⁾. The acid reaction of the final medium is inhibitory to a large number of bacteria making the media particularly useful for cultivating fungi and aciduric microorganisms. For isolation of fungi from contaminated specimens, a selective medium should be inoculated simultaneously. Incubate cultures for 4 to 6 weeks before reporting as negative.

Methodology

Suspend 45.0 grams of powder media in 1000 ml distilled water. Shake well & heat to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. DO NOT OVERHEAT. Mix well and pour in sterile Petri plates.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder.

Gelling

Firm, comparable with 1.5% Agar gel.

Colour and Clarity of prepared medium

Light amber coloured clear to slightly opalescent gel forms in Petri plates.

Reaction

Reaction of 4.5% w/v aqueous solution at 25°C. pH : 5.4±0.2

pH Range 5.20-5.60

Cultural Response/ characteristics

DM 2313: Cultural characteristics observed after an incubation at 25 - 30°C for upto 5 days.



Dehydrated Culture Media
Bases / Media Supplements

Organism	Inoculum (CFU)	Growth	Recovery
			>=70%
* <i>Aspergillus brasiliensis</i> ATCC 16404	50-100	good-luxuriant	
<i>Candida albicans</i> ATCC 10231	50-100	good-luxuriant	>=70%
<i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant(Inhibited on media with low pH)	>=70%
<i>Lactobacillus casei</i> ATCC 9595	50-100	good-luxuriant	>=70%
<i>Saccharomyces cerevisiae</i> ATCC 9763	50-100	good-luxuriant	>=70%
<i>Trichophyton rubrum</i> ATCC 28191	50-100	good-luxuriant	>=70%
<i>Penicillium notatum</i> ATCC 10108	50-100	good-luxuriant	>=70%
<i>Trichophyton gallinae</i> ATCC 22243	50-100	good-luxuriant	>=70%
<i>Trichophyton mentagrophytes</i> ATCC 9533	50-100	good-luxuriant	>=70%
<i>Trichophyton ajelloi</i> ATCC 24885	50-100	good-luxuriant	>=70%
*Key: Formerly known as <i>Aspergillus niger</i>			

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Carlier G. I. M., 1984, Brit. J. Derm. Syph., 60:61
2. Sabouraud R., 1892, Ann. Dermatol. Syphil. 3: 1061.
3. Merkblatt 18: Verpackgs- Rdsch, 1974, 25/1: Techn- Wiss. Beilage, 5-8
4. Merkblatt 19: Verpackgs- Rdsch, 1974, 25/6: Techn- Wiss. Beilage, 569-575
5. Merkblatt 21: Verpackgs- Rdsch, 1974, 25/7: Techn- Wiss. Beilage, 53-55
6. Murray P. R., Baron J. H., Pfaller M. A., Jorgensen J. H. and Tenover F. C., (Ed.), 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

