

## Technical Information

### Plate Count Agar with BCP

**Product Code: DM 2351**

**Application:** Plate Count Agar w/ BCP is used for enumeration of Lactobacilli in cultured milk, yoghurt and sour creams.

#### Composition\*\*

Ingredients	Gms / Litre
Peptic digest of animal tissue	5.000
Yeast extract	2.500
Dextrose	1.000
Polysorbate 80 (Tween 80)	1.000
L-Cysteine	0.100
Bromo cresol purple	0.040
Agar	15.000
Final pH ( at 25°C)	6.8±0.2

\*\*Formula adjusted, standardized to suit performance parameters

#### Principle & Interpretation

Plate Count Agar with BCP is a slight modification of the medium recommended by APHA <sup>(1)</sup> This medium is used for enumeration of Lactobacilli in cultured milk, yoghurt and sour creams. Lactobacilli grow on this medium and form colonies with yellowish peripheries in the depth and on the surface of the medium. Polysorbate 80 and L- cysteine enhanced the growth of lactobacilli on this medium. *Lactobacillus bulgaricus* invariably forms yellowish colonies that make its detection and determination very easy.

Peptic digest of animal tissue supply nitrogenous and carbonaceous compounds. Yeast extract provides vitamin B complex and dextrose is the fermentable carbohydrate and energy source. Bromocresol purple is a pH indicator. Plate Count Agar can be incubated at 30°C for upto 72 hours for detection of mesophilic organisms. Polysorbate 80 supplies fatty acids required for the metabolism of lactobacilli.

#### Methodology

Suspend 24.64 grams of powder media in 1000 ml. distilled water. Shake well & heat to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.

#### Quality Control

##### Physical Appearance

Light yellow to light green homogeneous free flowing powder

##### Gelling

Firm, comparable with 1.5% Agar gel

##### Colour and Clarity of prepared medium

Purple coloured clear to slightly opalescent gel forms in Petri plates

##### Reaction

Reaction of 2.46% w/v aqueous solution at 25°C. 6.8±0.2 pH : 6.8±0.2

**pH range** 6.60-7.00

##### Cultural Response/Characteristics

DM 2351: Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours.



Dehydrated Culture Media  
Bases / Media Supplements

Organism	Inoculum (CFU)	Growth
<i>Lactobacillus acidophilus</i> ATCC 4356	50-100	luxuriant
<i>Lactobacillus lactis</i> ATCC 19435	50-100	Luxuriant
<i>Lactobacillus bulgaricus</i> ATCC 11842	50-100	Luxuriant
<i>Streptococcus thermophilus</i> ATCC 14485	50-100	Luxuriant

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

## Further Reading

1. Downes F. P. and Ito K., (Eds.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., APHA, Washington, D.C.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

