

Technical Information

SD Agar

Product Code: DM 2371

Application: - SD Agar is used for the growth of yeasts for molecular biology purposes.

Composition**

Ingredients	Gms / Litre
Ammonium sulphate	5.000
Dextrose	20.000
Biotin	0.00002
Calcium pantothenate	0.002
Folic acid	0.000002
Inositol	0.010
Niacin	0.0004
p-Amino benzoic acid (PABA)	0.0002
Pyridoxine hydrochloride	0.0004
Riboflavin	0.0002
Thiamine hydrochloride	0.0004
Boric acid	0.0005
Copper sulphate	0.00004
Potassium iodide	0.0001
Ferric chloride	0.0002
Manganese sulphate	0.0004
Sodium molybdate	0.0002
Zinc sulphate	0.0004
Potassium dihydrogen phosphate	0.850
Dipotassium hydrogen phosphate	0.150
Magnesium sulphate	0.500
Sodium chloride	0.100
Calcium chloride	0.100
Agar	20.000
Final pH (at 25°C)	6.9±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

This medium is formulated as per Wickerham ⁽²⁾ for studying colonial and cellular morphology of yeasts. Dextrose provides on energy source, while the salts serve as a source of nutrition for metabolic reactions.

Methodology

Suspend 46.71 grams of powder media in 1000 ml distilled water. Shake well & heat to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.



Dehydrated Culture Media
Bases / Media Supplements

Quality Control

Physical Appearance

Cream to beige homogeneous free flowing powder.

Gelling

Firm, comparable with 2.0% agar gel.

Colour and Clarity of prepared medium

Light amber coloured, opalescent gel forms in precipitate.

Reaction

Reaction of 4.6% w/v aqueous solution at 25°C. pH 6.9 ± 0.2 pH : 6.9 ± 0.2

pH Range 6.70-7.10

Cultural Response/ characteristics

DM 2371: Cultural characteristics observed after an incubation at 35-30°C for 48-72 hour.

Organism	Inoculum (CFU)	Growth	Recovery
<i>Candida albicans</i> ATCC 10231	50-100	luxuriant	>=70%
<i>Saccharomyces cerevisiae</i> ATCC 9763	50-100	luxuriant	>=70%

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Sherman, F., 1991, Meths. Enzymol. 194:3
2. Wickerham, 1951, U.S. Dept. Agri. Tech. Bull. No. 1029.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

