

## Technical Information

### Fluid Selenite Cystine Broth (Twin Pack)

#### Product Code: DM 2533I

**Application:** - This medium is recommended as an enrichment medium for the isolation of Salmonellae from faeces, urine or other pathological materials.

#### Composition\*\*

Ingredients	Gms / Litre
Part A	-
Casein enzymic hydrolysate	5.000
Lactose	4.000
Disodium phosphate. 12H <sub>2</sub> O	10.000
L-Cystine	0.010
Part B	-
Sodium hydrogen selenite	4.000
Final pH ( at 25°C)	7.0±0.2

\*\*Formula adjusted, standardized to suit performance parameters

#### Principle & Interpretation

Selective inhibitory effects of selenite was first resported by Klett <sup>(1)</sup>. Guth <sup>(2)</sup> used this study to isolate *Salmonella* serotype Typhi. Leifson studied effects of selenite and formulated a medium. Fluid Selenite Cystine Medium is a modification of Leifson's <sup>(3)</sup> formula with added cystine <sup>(4)</sup>. The formulation corresponds to recommendations of AOAC <sup>(5)</sup> for the detection of Salmonellae in foodstuff, particularly egg products and included by APHA <sup>(6, 7)</sup>, USP <sup>(8)</sup>. This medium is recommended by ISO Committee <sup>(9)</sup> also as a selective enrichment medium for the detection of Salmonellae. 10 ml of culture from Buffered Peptone Water (DM1614S) is inoculated in Fluid Selenite Cystine Broth (DM2533I) and further sub-cultured on Brilliant Green Agar w/Phosphate (DM1971S).

Selenite Cystine Broth is useful to enhance the detection of low numbers of organisms from asymptomatic or convalescent patients <sup>(10)</sup> epidemiologically and for detecting *Salmonella* in the nonacute stages of illness when organisms are scanty in the faeces. Casein enzymic hydrolysate provides nitrogenous substances. Lactose maintains the pH in medium as selenite is reduced by bacterial growth and alkali is produced. An increase in pH lessens the toxicity of the selenite and results in overgrowth of other bacteria. The acid produced by bacteria due to lactose fermentation serves to maintain a neutral pH. Phosphate maintains a stable pH and also lessens the toxicity of selenite. L-cystine improves recovery of Salmonellae. Enriched broth is subcultured on solid medium. Do not incubate the broth longer than 24 hours as inhibitory effect of selenite reduces after 6 - 12 hours of incubation <sup>(11)</sup>.

#### Methodology

Suspend 4.0 grams of Part B in 1000 ml distilled water. Add 13.0 grams of dehydrated Part A medium. Warm to dissolve the medium completely. Distribute in sterile test tubes. Sterilize in a boiling water bath or in a free flowing steam for 10 minutes. DO NOT AUTOCLAVE. Excessive heating is detrimental. Discard the prepared medium if large amount of selenite is reduced. (Indicated by red precipitate at the bottom of tube/bottle).

Caution: Sodium hydrogen selenite (Sodium bi-selenite) is very toxic, corrosive agent and causes teratogenicity and hence should be handled with great care. Upon contact with skin, wash immediately with a lot of water.



Dehydrated Culture Media  
Bases / Media Supplements

## Quality Control

### Physical Appearance

Part A : Off-white to light yellow Part B : White to cream homogeneous free flowing powder

### Colour and Clarity of Prepared medium

Light yellow clear to slightly opalescent solution

### Reaction

Reaction of [(1.3% w/v) Part A and (0.4% w/v) Part B] at 25°C. pH : 7.0±0.2

pH range 6.80-7.20

### Cultural Response/ characteristics

DM2533I Cultural characteristics observed after an incubation at 35 - 37°C for 18 - 24 hours when subcultured on MacConkey Agar (DM1081).

Organism	Inoculum (CFU)	Recovery (increase in numbers)	Colour of colony
Escherichia coli ATCC 25922	50-100	little-none	pink w/ bile ppt
Salmonella Choleraesuis ATCC 12011	50-100	luxuriant	colourless
Salmonella Typhi ATCC 6539	50-100	luxuriant	colourless
Salmonella Typhimurium ATCC 14028	50-100	luxuriant	colourless

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

**Prepared Media:** 2-8° in sealable plastic bags for 2-5 days.

## Further Reading

1. Klett A., 1900, Zeitsch Fer Hyg. Und. Infekt, 33: 137.
2. Guth F., 1916, Zbl. Bakt. I. Orig., 77:487.
3. Leifson E., 1936, Am. J. Hyg., 24(2) : 423.
4. North W.R. and Bartram M.T., 1953, Appl. Microbiol., 1:130.
5. AOAC, 1978, Bacteriological Analytical Manual, 5th ed., AOAC, Washington, ,,DC.
6. Vanderzant C. and Splittstoesser D. (Eds.), 2001, Compendium of Methods for,, the Microbiological Examination of Foods, 4th ed., APHA, Washington, DC.
7. APHA, 2001, Standard Methods for the Examination of Dairy Products, 17th,, ed., APHA Inc., Washington, D.C.
8. United States Pharmacopeia, 2004, USP27/NF22, U. S. Pharmacopeial,, Convention, Inc., Rockville, MD.
9. International Organization for Standardization (ISO), 1993, Draft ISO/DIS,, 6579.
10. Kelly, Brenner and Farmer, 2003, Manual of Clinical Microbiology, 8th ed., ,,Lennett et al. (Eds.), ASM, Washington, D.C. 1
1. Chattopadhyay W. and Pilford J. N., 1976, Med. Lab. Sci., 33:191.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specificatons for identity and performens parameters.

