

Technical Information

B12 Culture Agar (L. leichmannii Maintenance Medium)

Product Code: DM 1035

Application: B12 Culture Agar is recommended for the propagation, cultivation and maintenance of *Lactobacillus leichmannii* in the Vitamin B12 Assay. Us 7830 used as the test organism

Composition**

Ingredients	Gms / Litre
Peptic digest of animal tissue	7.500
Yeast extract	7.500
Dextrose	10.000
Monopotassium phosphate	2.000
Tomato juice (from 100 ml)	5.000
Polysorbate 80	0.100
Agar	10.000
Final pH (at 25°C)	6.8±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

B12 Culture Agar recommended by USP for cultivation and maintenance of (*Lactobacillus delbrueckii subsp.lactis* ATCC 7830) is used as a test bacterium during the estimation of vitamin B12. ⁽¹⁾ *Lactobacillus* species have very exacting nutritional requirements for amino acids and vitamins. These make them to compete nutritionally in the environment. *Lactobacillus* species grow poorly on non-selective media. Kulp ⁽²⁾ found that the growth of *Lactobacillus acidophilus* was enhanced with the addition of tomato juice. This was also reported earlier by Mickle and Breed ⁽³⁾ Peptic digest of animal tissue serves as a source of nitrogen and amino acids. Yeast extract is the vitamin source. Tomato juice is added to create the proper acidic environment. Dextrose is the carbon source and Polysorbate 80 acts as an emulsifier. Monopotassium phosphate provides buffering capacity. Stock cultures of *Lactobacillus leichmannii* ATCC 7830 are prepared by stab inoculation of 3 or more tubes. These stab cultures are made at least 3 times in a week. Do not use the culture for preparing assay inoculum if it is over 4 days old. Before using a fresh culture for assay, make at least 10 successive transfers of the culture in 15 days period. Incubate the culture for 16-24 hours at 35°C but hold constant within 0.5°C. After incubation, store at 2-8°C.

Methodology

Suspend 42.1 grams of powder media in 1000 ml distilled water. Shake well and heat to dissolve the medium completely. Dispense in 10 ml amounts in tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool the tubed medium in an upright position with rapidity to avoid colour formation due to overheating.



Dehydrated Culture Media
Bases / Media Supplements

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.0% Agar gel.

Colour and Clarity of prepared medium

Medium amber coloured, clear to slightly opalescent gel forms in tubes.

Reaction

Reaction of 4.21% w/v aqueous solution at 25°C. pH : 6.8±0.2

pH Range 6.80-7.20

Cultural Response/Characteristics

DM1035: Cultural characteristics observed after an incubation at 35-37°C for 16-24 hours.

Organism	Inoculum (CFU)	Growth
<i>Lactobacillus leichmannii</i> ATCC 7830	50-100	good-luxuriant

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8⁰ in sealable plastic bags for 2-5 days.

Further Reading

1. The United States Pharmacopoeia, 2006, USP 29/ NF 24, The United States Pharmacopoeial Convention, Rockville, MD.
2. Kulp and White, 1932, Science 76:17.
3. Mickle and Breed, 1925, Technical Bulletin 110, NY State Agriculture Ex. station, Geneva, N.Y.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

