

Technical Information

Urea Agar Base (Christensen)

Product Code: DM 1112S

Application: Urea Agar Base is recommended for the detection of urease production, particularly by *Proteus vulgaris*, Micrococci and paracolon organisms.

Composition**

Ingredients	Gms / Litre
Dextrose	1.000
Peptic digest of animal tissue	1.500
Sodium chloride	5.000
Monopotassium phosphate	2.000
Phenol red	0.012
Agar	15.000
Final pH (at 25°C)	6.8±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Urea Agar Base Media is a slight modification of Christensen formulation ^(1,2) and is recommended by BIS ^(3,4) for identification of urease activity. Rustigian and Stuart ⁽⁵⁾ had originally devised a medium to detect urease activity. These media differentiate between rapid urease positive *Proteus species* and other urease positive organisms like *Citrobacter*, *Enterobacter* and *Klebsiella* and the bacteria other than *Enterobacteriaceae*. Christensen observed that addition of peptic digest of animal tissue, dextrose and reduced content of buffer helps in an early luxuriant growth of the microbes.

Heavy inoculum of growth is inoculated on the surface of the slants. When urea is utilized, ammonia is produced during incubation which makes the medium alkaline, showing a pink-red colour by the change in the phenol red indicator. Prolonged incubation may cause alkaline reaction in the medium. As negative control medium without urea should be included in test proper.

Methodology

Suspend 24.51 grams of powder media in 950 ml distilled water. Shake well & heat to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 50°C and aseptically add 50 ml of sterile 40% Urea Solution (MS2048) and mix well. Dispense into sterile tubes and allow to set in the slanting position. Do not overheat or reheat the medium as urea is heat.

Quality Control

Physical Appearance

Light pink coloured homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Yellowish orange coloured clear gel forms in tubes as slants.

Reaction

Reaction of 2.45% w/v aqueous solution at 25°C. pH : 6.8±0.2

pH range 6.60-7.00

Cultural Response/Characteristics

DM 1112S: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth	Urease
<i>Enterobacter aerogenes</i> ATCC 13048	50-100	luxuriant	Negative reaction, no change
<i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	Negative reaction, no change
<i>Klebsiella pneumoniae</i> ATCC 13883	50-100	luxuriant	Positive reaction, cerise colour
<i>Proteus mirabilis</i> ATCC 12453	50-100	luxuriant	Positive reaction, cerise colour
<i>Proteus vulgaris</i> ATCC 13315	50-100	luxuriant	Positive reaction, cerise colour
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	luxuriant	Negative reaction, no change

Storage and Shelf Life**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.**Prepared Media:** 2-8⁰ in sealable plastic bags for 2-5 days**Further Reading**

1. Christensen, W.B., 1946, J. Bact., 52:461.
2. MacFaddin J., 1980, Biochemical Tests for Identification of Medical Bacteria, 2nd ed., Williams and Wilkins, Baltimore.
3. Bureau of Indian Standards, IS : 5887 (Part I) - 1976, reaffirmed 1986.
4. Bureau of Indian Standards, IS : 5887 (Part III) - 1999.
5. Rustigian and Stuart, 1941, Proc. Soc. Exp. Biol. Med., 47:108.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at CDH is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.