

Technical Information

Dextrose Starch Agar

Product Code: DM 1183

Application: Dextrose Starch Agar is used for propagating pure cultures of *Neisseria gonorrhoeae* and other fastidious organisms.

Composition**

Ingredients	Gms / Litre
Proteose peptone	15.000
Dextrose	2.000
Starch, soluble	10.000
Sodium chloride	5.000
Disodium phosphate	3.000
Gelatin	20.000
Agar	10.000
Final pH (at 25°C)	7.3±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Neisseria meningitidis, the causative agent of meningitis, is responsible for a large amount of morbidity and mortality throughout the world while *Neisseria gonorrhoeae* is the causative agent of the sexually transmitted disease gonorrhea that rank second to chlamydia (CDC). These fastidious organisms can be cultivated on Dextrose Starch Agar. The medium is highly nutritious and supports the luxuriant growth of various fastidious organisms like *N. meningitidis*, *Streptococcus pyogenes* and *Streptococcus pneumoniae* without adding any supplements. Organisms lacking the ability of starch hydrolysis can be maintained on this medium. Organism capable of hydrolyzing starch will create acidic conditions thereby making it unsuitable for maintenance. Dextrose Starch Agar was used to test the activity of various antibiotics against *Neisseria* species by the agar dilution technique as demonstrated by Wilkins, Lewis and Barbiers⁽¹⁾. *N. meningitidis* grow luxuriantly on this medium, when the plates are kept in 4-6% CO₂ environment or in the presence of abundant moisture. Swancara⁽²⁾ has described a method of obtaining partial carbon-dioxide tension and this can be used for incubation of Dextrose Starch Agar plates inoculated with *N. meningitidis*.

Proteose peptone and gelatin serve as sources of nitrogen and carbon essential for microbial growth. Dextrose serves as the energy source. Starch neutralizes toxic fatty acids that may be present in the agar. Sodium chloride maintains the osmotic balance and buffering is achieved by inclusion of disodium phosphate.

Dextrose Starch Agar prepared in half strength is a good medium for maintaining stock cultures of gonococci. The medium normally contains a flocculent precipitate, which does not affect the nutritive value of the medium. It is necessary to have the incubation atmosphere saturated with moisture while cultivating gonococci. Suitable conditions can be achieved, if the plates are incubated in a closed container containing cotton or towel saturated with water. Best results are obtained on a solid medium with a moist surface.

Methodology

Suspend 65 grams of powder media in 1000 ml distilled water. Shake well and heat to dissolve the medium completely. Dispense in tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool the tubed medium in a slanted position.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.0% Agar gel and 2.0% gelatin.

Colour and Clarity of prepared medium

Light amber coloured, opalescent gel with flocculent precipitate forms in tubes as slants

Reaction

Reaction of 6.5% w/v aqueous solution at 25°C. pH : 7.3±0.2

pH Range 7.10-7.50

Cultural Response/Characteristics

DM1183: Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours in an anaerobic conditions.

Organism	Inoculum (CFU)	Growth
<i>Neisseria gonorrhoeae</i> ATCC 19424	50-100	luxuriant
<i>Neisseria meningitidis</i> ATCC 13090	50-100	luxuriant
<i>Streptococcus pneumoniae</i> ATCC 6303	50-100	luxuriant
<i>Streptococcus pyogenes</i> ATCC 19615	50-100	luxuriant

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Wilkins, Lewis and Barbiers, 1956, Antibiot. Chemother., 6:149.
2. Swancara, 1948, Am. J. Med. Tech., 14:214.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.