

## Technical Information

### Dey-Engley Neutralizing Broth Base

#### Product Code: DM 1187

**Application:** Dey-Engley Neutralizing Broth Base is used in disinfectant testing where neutralization of the antiseptics and disinfectants is important for determining its bactericidal activity.

#### Composition\*\*

Ingredients	Gms / Litre
Casein enzymic hydrolysate	5.000
Yeast extract	2.500
Dextrose	10.000
Bromocresol purple	0.020
Final pH ( at 25°C)	7.6±0.2

\*\*Formula adjusted, standardized to suit performance parameters

#### Principle & Interpretation

Dey Dey-Engley Neutralizing Broth neutralizes broad spectrum antiseptics and disinfectants including quaternary ammonium compounds, phenolics, iodine and chlorine preparations, mercurials, formaldehyde and glutaraldehyde. This medium is used for the neutralization and testing of antiseptics and disinfectants according to the procedure of Engley and Dey<sup>(1)</sup> who formulated this media. A strongly bacteriostatic substance inhibits the growth and reproduction of bacteria without killing them. These bacteria have the ability to cause infection under favorable conditions. Dey -Engley Neutralizing Broth Base does not contain the neutralizing components.

Casein enzymic hydrolysate provides essential nutrients. Dextrose is an energy source. Yeast extract is also a rich source of vitamin B-complex.<sup>(1)</sup> Bromocresol purple is an indicator for dextrose utilization. Therefore, bromo cresol purple and dextrose are added to the medium. Those organisms that ferment dextrose will turn the medium from purple to yellow.<sup>(1)</sup>

##### Neutralization Test :

For testing disinfectants, prepare two sets of test tubes, one containing 9 ml Dey-Engley Neutralizing Broth (DM2062) and other with 9 ml Dey-Engley Neutralizing Broth Base (DM1187). Add 1 ml of disinfectant under test. Mix well and allow it to stand for 15 minutes. Inoculate 0.1 ml of 1:100,000 dilution of overnight broth cultures and incubate at 37°C for 48 hours. Growth is indicated by a colour change from purple to yellow or pellicle formation. Growth in Neutralizing Broth and no growth in Neutralizing Broth Base indicate neutralization of disinfectant. To check bactericidal activity, both broth tubes are inoculated on D/E Neutralizing Agar (DM1186). Positive growth from negative tubes of Neutralizing Broth Base indicates bacteriostatic substance while negative growth indicates a bactericidal disinfectant. All positive tubes should show growth on Dey-Engley Neutralizing Agar. The control disinfectants used in test procedure are 2% chlorine, 2% formaldehyde, 1% glutaraldehyde, 2% iodine, 2% phenol, 1/750 quaternary ammonium compounds, 1/1000 mercurials etc. Quality Control :

#### Methodology

Suspend 17.52 grams of powder media in 1000 ml distilled water. Shake well and heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and dispense as desired.

#### Quality Control

##### Physical Appearance

Light yellow to bluish grey homogeneous free flowing powder

##### Colour and Clarity of prepared medium

Purple coloured, opalescent solution in tubes.

##### Reaction

Reaction of 1.75% w/v aqueous solutions at 25°C. pH : 7.6±0.2



Dehydrated Culture Media  
Bases / Media Supplements

#### pH Range

7.40-7.80

#### Cultural Response/Characteristics

DM1187: Cultural characteristics observed after an incubation at 35-37°C for 40-48 hours.

Organism	Inoculum (CFU)	Growth
<i>Bacillus subtilis</i> ATCC 6633	50-100	Luxuriant
<i>Escherichia coli</i> ATCC 25922	50-100	Luxuriant
<i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	Luxuriant
<i>Salmonella Typhimurium</i> ATCC 14028	50-100	Luxuriant
<i>Staphylococcus aureus</i> ATCC 25923	50-100	luxuriant

### Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

**Prepared Media:** 2-8<sup>0</sup> in sealable plastic bags for 2-5 days.

### Further Reading

1. Engley and Dey, 1970. Chem. Spec. Manuf. Assoc. Proc., Mid-Year Meet., p. 100.

### Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

