

Technical Information

NIH Agar

Product Code: DM 1194

Application: NIH Agar is used for sterility testing and for the cultivation and maintenance of isolates from sterility testing of biological products.

Composition**

Ingredients	Gms / Litre
Casein enzymic hydrolysate	15.000
Yeast extract	5.000
Dextrose	5.500
Sodium chloride	2.500
L-Cystine	0.050
Agar	15.000
Final pH (at 25°C)	7.1±0.2

** Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

This medium can be used for sterility testing and also for cultivating the isolates from biological products tested for sterility. This medium is also recommended by the National Institute of Health (NIH) for sterility testing of biological products of turbid nature ⁽³⁾. NIH Medium has a similar composition as Fluid Thioglycollate Medium, except sodium thioglycollate has resazurin and agar concentration is more in NIH Medium than in Fluid Thioglycollate Medium NIH Agar is formulated according to the agar medium specified by USPHS sterility test ⁽¹⁾.

NIH medium is a nutritious medium containing nutrients like casein enzymic hydrolysate, yeast extract and the amino acid L-cystine. It contains the fermentable carbohydrate dextrose and sodium chloride for maintaining osmotic equilibrium. NIH Medium is devoid of sodium thioglycollate. U.S. Pharmacopoeia ⁽²⁾ has recommended using this medium with sodium thioglycollate (0.05%) or thioglycollic acid (0.03%) for the sterility testing of biological products containing mercurial preservatives, since sodium thioglycollate neutralizes the bacteriostatic effect of mercuric compounds ^(4,5).

Methodology

Suspend 43.05 grams of powder media in 1000 ml distilled water. Shake well and heat to dissolve the medium completely. Dispense into test tubes or flasks as desired. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.5% Agar gel

Colour and Clarity of prepared medium

Light amber coloured clear to slightly opalescent gel forms in Petri plates

Reaction

Reaction of 4.3% w/v aqueous solution at 25°C. pH : 7.1±0.2

pH Range 6.90-7.30

Cultural Response/ characteristics

DM 1194: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours and with addition of sodium thioglycollate.



Dehydrated Culture Media
Bases / Media Supplements

Organism	Inoculum (CFU)	Growth	Recovery
Escherichia coli ATCC 25922	50-100	good-luxuriant	>=70%
Streptococcus mitis ATCC 9895	50-100	good-luxuriant	>=70%
Staphylococcus aureus ATCC 25923	50-100	good-luxuriant	>=50%
Streptococcus pyogenes ATCC 19615	50-100	good-luxuriant	>=70%
Bacillus subtilis ATCC 6633	50-100	good-luxuriant	>=50%
Bacteroides vulgatus ATCC 8482	50-100	good-luxuriant	>=50%
Candida albi cans ATCC 10231	50-100	good-luxuriant	>=50%
Micrococcus luteus ATCC 9341	50-100	good-luxuriant	>=50%
Clostridium sporogenes ATCC 11437	50-100	good-luxuriant	>=50%

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. USPHS Reg., 73, 730: Federal Register, 1970, Vol. 35, No. 0171, p. 13:930.
2. The United States Pharmacopoeia, 2006, USP29/NF24. The United States Pharmacopoeial Convention, Rockville, MD.
3. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams and Wilkins, Baltimore
4. Nungester, Hood and Warren, 1943, Proc. Soc. Exp. Biol. Med. 52 : 287
5. Portwood, 1944, J. Bacteriol., 48: 255.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conforms solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate.
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

