

Technical Information

Lactobacilli Agar, AOAC

Product Code: DM 1366

Application: Lactobacilli Agar, AOAC is used for carrying the stock cultures used in microbiological assays of the B-vitamins.

Composition**

Ingredients	Gms / Litre
Peptonized milk	15.000
Yeast extract	5.000
Dextrose	10.000
Tomato juice (100 ml)	5.000
Monopotassium phosphate	2.000
Polysorbate 80	1.000
Agar	10.000
Final pH (at 25°C)	6.8±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Lactobacilli Agar, AOAC was devised by Loy ⁽¹⁾ and recommended by AOAC ⁽²⁾ for Microbiological assay of Vitamin B is used for the cultivation and maintenance of stock cultures of *Lactobacillus casei* ATCC 7469, *Lactobacillus fermentum* ATCC 9338, *Lactobacillus leichmannii* ATCC 4797, and *Lactobacillus viridescens* ATCC 12706. These cultures are grown by stab inoculation of sterile tubed medium followed by incubation for 18-24 hours at a constant temperature between 30-40°C. The cultures are then stored at 2-8°C.

Peptonized milk and yeast extract provide essential growth nutrients. Dextrose is the energy source. Phosphate provides buffering system while tomato juice helps in lowering the pH. Polysorbate 80 serves as an emulsifier.

Before using a culture in any assay, at least 2 successive transfers during a 1-2 week period are essential. Any culture older than one week should not be used.

Methodology

Suspend 48 grams of powder media in 1000 ml distilled water. Shake well & heat to dissolve the medium completely. Distribute into tubes in 10 ml amounts and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool the tubed agar medium in an upright position.

Quality Control

Physical Appearance

Cream to yellow homogeneous free flowing powder

Gelling

Firm, comparable with 1.0% Agar gel.

Colour and Clarity of prepared medium

Medium amber coloured clear to slightly opalescent gel forms in tubes as butts

Reaction

Reaction of 4.8% w/v aqueous solution at 25°C. pH : 6.8±0.2

pH Range:- 6.60-7.00

Cultural Response/Characteristics

DM 1366: Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours



Dehydrated Culture Media
Bases / Media Supplements

Organism	Inoculum (CFU)	Growth
<i>Enterococcus hirae</i> ATCC 8043	50-100	luxuriant
<i>Lactobacillus casei</i> ATCC 7469	50-100	luxuriant
<i>Lactobacillus leichmannii</i> ATCC 7830	50-100	luxuriant
<i>Lactobacillus plantarum</i> ATCC 8014	50-100	luxuriant

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8⁰ in sealable plastic bags for 2-5 days.

Further Reading

1. Loy, 1958, J. AOAC, 4:61.
2. Williams, (Ed.), 2005, Official Methods of Analysis of the Association of Official Analytical Chemists, 19th Ed., AOAC, Washington, D.C

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performance parameters.

