

Technical Information

Lysine Decarboxylase Broth without Peptone

Product Code: DM 1376I

Application: Lysine Decarboxylase Broth w/o Peptone are used for differentiating *Salmonella* Arizonae from the Bethesda Ballerup group of *Enterobacteriaceae*

Composition**

Ingredients	Gms / Litre
L-Lysine hydrochloride	5.000
Yeast extract	3.000
Dextrose	1.000
Bromocresol purple	0.015
Final pH (at 25°C)	6.8±0.2

**Formula adjusted, standardized to suit performance parameters

Principal & Interpretation

Decarboxylase media were first described by Moeller ⁽¹⁻³⁾ for detecting lysine and ornithine decarboxylase and arginine dihydrolase. Falkow developed a lysine decarboxylase medium for the identification and differentiation of *Salmonella* and *Shigella* ⁽⁴⁾. Falkow's Medium was further modified by Taylor ⁽⁵⁾ by removing peptone from the formulation (DM1376I), thus eliminating false positives caused by *Citrobacter freundii* and its paracolons. Taylor's modification has same advantage of Falkow's formulation over Moeller; it does not require the special conditions of anaerobic culture and low pH.

During the initial stages of incubation, fermentation of dextrose by the organisms, with acid production results in a colour change of the indicator to yellow. On further incubation, if L-Lysine is decarboxylated to cadaverine, there will be an alkaline reaction and the indicator colour will then revert back to purple. If the colour remains yellow, the decarboxylase reaction is negative.

Yeast extract provide essential growth nutrients. Dextrose is the fermentable carbohydrate and bromo cresol purple is the pH indicator. Dextrose non-utilizers will not show any change in the medium colour. Use light inocula and do not read the tests under 24 hours incubation as some organisms require longer incubation time of upto 4 days.

Inoculate 25 grams of the test sample into Buffered Peptone Water (DM1614S). After incubation at 35-37°C for 16-20 hours, inoculate into RVS Broth (DM2491) and Fluid Selenite Cystine Broth (DM2533I) and incubate at 35-37°C for 24-48 hours. From the second enrichment, streak a loopful on Brilliant Green Agar Base w/ phosphates (DM1971S). Presumptive *Salmonella* so isolated on (DM1971S) are further confirmed by performing biochemical testing using the following media i.e. Nutrient Agar, pH 7.0 (DM1561A), Triple Sugar Iron Agar (DM1021S), Urea Agar Base, Christensen (DM1112I), Lysine Decarboxylase Broth w/o peptone (DM1376I), VP test, Indole test.

Methodology

Suspend 9.01 grams in 1000 ml distilled water. Heat if necessary to dissolve the medium completely. Dispense 5 ml amount into screw-capped test tubes. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

Quality Control

Physical Appearance

Light yellow to greenish yellow homogeneous free flowing powder

Colour and Clarity of prepared medium

Purple coloured clear solution without any precipitate

Reaction

Reaction of 0.9% w/v aqueous solution at 25°C. pH : 6.8±0.2

pH Range 6.60-7.00

Cultural Response/Characteristics

DM 1376I: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Lysine decarboxylation
<i>Citrobacter freundii</i> ATCC 8090	50-100	Variable reaction
<i>Escherichia coli</i> ATCC 25922	50-100	Variable reaction
<i>Enterobacter aerogenes</i> ATCC 13048	50-100	Positive reaction, purple colour
<i>Klebsiella pneumoniae</i> ATCC 13883	50-100	Positive reaction, purple colour
<i>Proteus mirabilis</i> ATCC 25933	50-100	Negative reaction, yellow colour
<i>Proteus vulgaris</i> ATCC 13315	50-100	Negative reaction, yellow colour
<i>Salmonella Arizonae</i> ATCC 13314	50-100n	Positive reaction, purple colour
<i>Salmonella Paratyphi A</i> ATCC 9150	50-100	Negative reaction, yellow colour
<i>Salmonella Typhi</i> ATCC 6539	50-100	Positive reaction, purple colour
<i>Serratia marcescens</i> ATCC 8100	50-100	Positive reaction, purple colour
<i>Shigella dysenteriae</i> ATCC 13313	50-100	Negative reaction, yellow colour

Storage and Shelf Life**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.**Prepared Media:** 2-8^o in sealable plastic bags for 2-5 days.**Further Reading**

1. Moeller V., 1954, Acta. Pathol. Microbiol. Scand., 34:102.
 2. Moeller V., 1954, Acta. Pathol. Microbiol. Scand., 34:259.
 3. Moeller V., 1955, Acta. Pathol. Microbiol. Scand., 36:158.
 4. Falkow, 1958, Am. J. Clin. Pathol., 29:598.
- Taylor W. I., 1961, Appl. Microbiol., 9:487.

Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Do not use the products if it fails to meet specifications for identity and performance parameters.