

Technical Information

Lysine Medium Base

Product Code: DM 1642

Application: - Lysine Medium Base is used for isolation and enumeration of wild yeasts in pitching yeasts.

Composition**

Ingredients	Gms / Litre
Dextrose	44.500
Monopotassium phosphate	1.780
Magnesium sulphate	0.890
Calcium chloride	0.178
Sodium chloride	0.089
Adenine	0.00178
DL-Methionine	0.000891
L-Histidine	0.000891
DL-Tryptophan	0.000891
Boric acid	0.0000089
Zinc sulphate	0.0000356
Ammonium molybdate	0.0000178
Manganese sulphate	0.0000356
Ferrous sulphate	0.0002225
L-Lysine	1.000
Inositol	0.020
Calcium pantothenate	0.002
Aneurine	0.0004
Pyridoxine	0.0004
p-Amino benzoic acid (PABA)	0.0002
Nicotinic acid	0.0004
Riboflavin	0.0002
Biotin	0.000002
Folic acid	0.000001
Agar	17.800
Final pH (at 25°C)	5.0±0.2

**Formula adjusted, standardized to suit performance parameters

Principle & Interpretation

Walters and Thiselton⁽²⁾ used a liquid synthetic medium containing lysine as sole nitrogen source and found that many types of yeast utilize lysine. Later Morris and Eddy⁽¹⁾ also formulated solid lysine medium. Most of the Saccharomyces strains used in the brewery industry and other fermentative industries do not use lysine, whereas the wild type strains do. This property of lysine medium base help in the separation of both types of yeast.

The medium contains vitamins and trace elements, which is necessary to support metabolic activities of yeast. Lysine acts as the sole source of nitrogen, which is utilized by many types of yeast. Morris and Eddy⁽¹⁾ recommended surface inoculation of washed aliquots from the yeast mass; 0.2 ml suspension of 10⁷ cells/ml is the best. Sample is incubated at 25°C and examined daily, enumerating all the colonies that have grown (lysine positive). The degree of contamination is expressed as the number of wild yeast cells per million cells of the original inoculum. The number of cells in the inoculum is important as small number of cells about 100 to 1000 grow to a limited extent while 10,000 brewing yeast cells provide a direct measure of contaminant wild yeasts⁽³⁾.



Dehydrated Culture Media
Bases / Media Supplements

Methodology

Suspend 6.62 grams of powder media in 100 ml distilled water containing 1 ml of 50% potassium lactate (MS2123). Shake well & heat to boiling to dissolve the medium completely. DO NOT AUTOCLAVE. Cool to 50°C, adjust pH to 5.0 with 10% lactic acid and pour into sterile Petri plates.

Quality Control

Physical Appearance

White to cream homogeneous free flowing powder

Gelling

Firm, comparable with 1.78% Agar gel.

Colour and Clarity of prepared medium

Colourless to pale yellow clear to slightly opalescent opalescent gel forms in Petri plates

Reaction

Reaction of 6.6% w/v aqueous solution at 25°C. pH : 5.0±0.2

pH range

4.80-5.20

Cultural Response/Characteristics

DM 1642: Cultural characteristics observed after an incubation at 25-30°C upto 7 days.

Organism

Pichia fermentans ATCC 10651

Growth

Luxuriant

Storage and Shelf Life

Dried Media: Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

Prepared Media: 2-8° in sealable plastic bags for 2-5 days.

Further Reading

1. Morris E. O. and Eddy A. A, 1957, J. Inst. Brew. 63(1): 34.
2. Walters L. S. and Thiselton M. R., 1953, J. Inst. Brew. 59:401.
3. Fowell R. R., 1965, J. Appl. Bacteriol., 28:373.

Disclaimer :

User must ensure suitability of the product(s) in their application prior to use.

The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried a at **CDH** is true and accurate

Central Drug House Pvt. Ltd. reserves the right to make changes to specifications and information related to the products at any time.

Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.

Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents. Donot use the products if it fails to meet specifacatons for identity and performens parameters.

