

## Technical Information

### Dextrose Peptone Agar

**Product Code: DM 1649**

**Application:** - Dextrose Peptone Agar is recommended for the general cultivation of organisms.

### Composition\*\*

Ingredients	Gms / Litre
Peptic digest of animal tissue	20.000
Dextrose	10.000
Sodium chloride	5.000
Agar	15.000
Final pH ( at 25°C)	7.2±0.2

\*\*Formula adjusted, standardized to suit performance parameters

### Principle & Interpretation

Dextrose Peptone Agar is devised by Williams <sup>(1)</sup> for the cultivation of microorganisms, which are fastidious, or present in small numbers, and also for the enumeration of the thermophilic bacteria responsible for flat sour spoilage of canned foods. This medium is recommended by AOAC for the routine cultivation purpose <sup>(2)</sup>. Peptic digest of animal tissue supplies amino acids, peptides etc. for the growth of the organisms. Dextrose is the readily available energy source for the most of the organisms. The agar medium is also used as an excellent basal agar for the Glucose Blood Agar preparation. In the special Petri plates, it can support good growth of the anaerobic microorganisms.

### Methodology

Suspend 50 grams of powder media in 1000 ml distilled water. Shake well and heat, to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.

### Quality Control

#### Physical Appearance

Cream to yellow homogeneous free flowing powder

#### Gelling

Firm, comparable with 1.5% Agar gel

#### Colour and Clarity of prepared medium

Light yellow coloured, clear to slightly opalescent gel forms in Petri plates

#### Reaction

Reaction of 5.0% w/v aqueous solution at 25°C. pH : 7.2±0.2

#### pH Range 7.00-7.40

#### Cultural Response/Characteristics

DM1649: Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours.

Organism	Inoculum (CFU)	Growth	Recovery
<i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	>=70%
<i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	luxuriant	>=70%
<i>Staphylococcus aureus</i> ATCC 25923	50-100	luxuriant	>=70%
<i>Streptococcus pyogenes</i> ATCC 19615	50-100	luxuriant	>=70%



Dehydrated Culture Media  
Bases / Media Supplements

## Storage and Shelf Life

**Dried Media:** Store below 30°C in tightly closed container and use before expiry date as mentioned on the label.

**Prepared Media:** 2-8<sup>0</sup> in sealable plastic bags for 2-5 days.

## Further Reading

1. Williams O.B., 1936, Food Res., 1(3):217.
2. Association of Official Analytical Chemists, 1978, Bacteriological Analytical Manual, 5th ed., AOAC, Washington, D.C.

## Disclaimer :

- User must ensure suitability of the product(s) in their application prior to use.
- The product conform solely to the technical information provided in this booklet and to the best of knowledge research and development work carried at **CDH** is true and accurate
- **Central Drug House Pvt. Ltd.** reserves the right to make changes to specifications and information related to the products at any time.
- Products are not intended for human or animal diagnostic or therapeutic use but for laboratory, research or further manufacturing of diagnostic reagents extra.
- Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.
- Do not use the products if it fails to meet specifications for identity and performs parameters.

